

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : A61K 38/44, 48/00, A61P 9/10 // A61K 38/18, 31/555	A2	(11) International Publication Number: WO 00/12118 (43) International Publication Date: 9 March 2000 (09.03.00)
(21) International Application Number: PCT/US99/19823 (22) International Filing Date: 27 August 1999 (27.08.99) (30) Priority Data: 60/098,377 28 August 1998 (28.08.98) US 60/121,946 25 February 1999 (25.02.99) US (71) Applicant: PRESIDENT AND FELLOWS OF HARVARD COLLEGE [US/US]; 17 Quincy Street, Cambridge, MA 02138 (US). (71)(72) Applicants and Inventors: LEE, Mu-En [CN/US]; 102 Nardell Road, Newton, MA 02159 (US). PERRELLA, Mark, A. [US/US]; 33 Pond Avenue, #420, Brookline, MA 02146 (US). YET, Shaw-Fang [CN/US]; 9 Donald Circle, Andover, MA 01810 (US). (74) Agent: BEATTIE, Ingrid, A.; Fish & Richardson P.C., 225 Franklin Street, Boston, MA 02110-2804 (US).		(81) Designated States: AU, CA, IL, JP, NO, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: INHIBITING CARDIOMYOCYTE DEATH (57) Abstract The invention features methods of inhibiting cardiomyocyte death in a mammal by administering to the myocardium of the mammal a heme oxygenase (HO). Methods of preserving an organ for transplantation and methods of inhibiting vascular stenosis are also within the invention.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

- 1 -

INHIBITING CARDIOMYOCYTE DEATHRelated Application Information

5 This application claims priority from provisional application no. 60/121,946, filed on February 25, 1999, and provisional application no. 60/098,377, filed on August 28, 1998.

Statement as to Federally Sponsored Research

10 This invention was made with U.S. Government support under National Institutes of Health grants R01 GM53249, KO8 HL03274, and KO8 HL03194. The government has certain rights in the invention.

Background of the Invention

15 The invention relates to treatment of cardiovascular disease.

 Myocardial infarction is one of the most common diagnoses of hospitalized patients in western countries. In the United States, over 1.5 million myocardial
20 infarctions occur annually, and mortality from acute myocardial infarction is approximately 25 per cent. Thrombolytic therapy and reperfusion of ischemic myocardium, e.g., using percutaneous transluminal coronary angioplasty (PTCA), have decreased mortality
25 rates among patients who have suffered an acute myocardial infarction. Nevertheless, myocardial damage and heart failure resulting from a failure to achieve early revascularization or from reperfusion injury remain a significant clinical problem.

30 Summary of the Invention

 The invention features methods of minimizing myocardial damage by salvaging hypoxic myocardial tissue before it becomes irreversibly injured. For example, a method of inhibiting cardiomyocyte death in a mammal,
35 e.g., a human, who has suffered a myocardial infarction or who has myocarditis is carried out by locally

- 2 -

administering to the myocardium of the mammal a heme oxygenase (HO) polypeptide. Preferably, the HO polypeptide has the amino acid sequence of a naturally-occurring heme oxygenase-1 (HO-1), heme oxygenase-2 (HO-2), or heme oxygenase-3 (HO-3), or a biologically active fragment thereof.

Compositions, such as hemin, hemoglobin, or heavy metals, e.g., tin or nickel, that increase production of endogenous HO, are also administered to inhibit cardiomyocyte death or damage. For example, overexpression of HO-1 is induced in vascular cells by exposure to heme, heavy metals, endotoxin and hyperoxia, hyperthermia, shear stress and strain, UV light, or reactive oxygen species. By "overexpression" is meant a level of protein production that at least 20% greater than that present in the tissue under normal physiologic conditions. Preferably, the level of HO-1 expression in vascular tissue in the presence of an inducing agent is at least 20% greater than that in the absence of the inducing agent; more preferably, the level of expression is at least 50% greater, more preferably, the level of expression is at least 100% greater, and most preferably, the level of expression is at least 200% greater than that in the absence of an inducing agent. Inhibition of cardiomyocyte death is also achieved by locally administering to the myocardium of a mammal a DNA encoding a HO. HO expression by target cell, e.g., VSMC, is increased by administering to the cells exogenous DNA encoding HO, e.g., a plasmid containing DNA encoding human HO-1 or HO-2 under the control of a strong constitutive promoter. Oxidative stress leads to cell death by apoptosis and/or necrosis. By neutralizing reactive oxygen species, HO reduces cardiomyocyte damage and death due to oxidative stress.

- 3 -

The invention also includes a method of inhibiting cardiomyocyte death *in vitro* by contacting cardiomyocytes with an HO or DNA encoding an HO. For example, a method of preserving isolated myocardial tissue, e.g., a donor heart to be used for transplantation, is carried out by bathing or perfusing the tissue with a solution containing an HO or a DNA encoding an HO. The method allows prolonged storage of organs after removal from the donor and prior to transplantation into a recipient by reducing irreversible ischemic tissue damage. By "isolated myocardial tissue" is meant tissue that has been removed from a living or recently deceased mammal. Preferably, a donor heart is preserved in an HO solution for 0.5-6 hours prior to transplantation. More preferably, the organ is preserved for greater than 6 hours, e.g., 8, 10, 12, and up to 24 hours.

A method of inhibiting vascular stenosis or restenosis in a mammal, e.g., a human, is also within the invention. The method is carried out by locally administering to the site of a vascular injury or a site which is at risk of developing a stenotic lesion a compound which inhibits expression of HO-1, and as a result, VSMC proliferation. To inhibit vascular stenosis or restenosis (which occurs at a relatively late stage after the occurrence of a vascular injury), the compound is administered at least one month after an injury such as surgery or angioplasty. For example, such treatment is administered 3 weeks to several months (e.g., 2 months or 3 months) post-injury. Preferably, the compound inhibits transcription of the gene encoding HO-1 or inhibits translation of HO-1 mRNA into an HO-1 polypeptide in a vascular cell, e.g., a vascular smooth muscle cell (VSMC), of the mammal. The vascular cell is preferably an aortic smooth muscle cell, e.g., an aortic smooth muscle cell located in the region of an artery

- 4 -

affected by vascular stenosis or restenosis such as the site of balloon angioplasty or coronary bypass surgery. For example, transforming growth factor- β 1 (TGF- β 1) is administered to inhibit production of HO-1 mRNA and HO gene product.

For antisense therapy, the compound is a antisense nucleic acid molecule containing at least 10 nucleotides, the sequence of which is complementary to an mRNA encoding all or part of a wild type HO polypeptide.

10 Preferably, the compound, e.g., an antisense oligonucleotide or antisense RNA produced from an antisense template, inhibits HO expression. The antisense nucleic acid inhibits HO expression by inhibiting translation of HO mRNA. For example,

15 antisense therapy is carried out by administering a single stranded nucleic acid complementary at least a portion of HO mRNA to interfere with the translation of mRNA into protein, thus reducing the amount of functional HO produced in the cell. The method includes the step of

20 identifying a mammal having undesired vascular stenosis or restenosis or at risk of developing such a condition. For example, the mammal to be treated is one who needs or has recently undergone PTCA, coronary artery bypass surgery, other vascular injury, that stimulates vascular

25 smooth muscle cell proliferation that results in undesired vascular stenosis or restenosis.

For treatment of a vascular injury soon after the injury or stress has occurred, an HO polypeptide, e.g., HO-1, or a nucleic acid encoding an HO polypeptide, is

30 administered to a mammal within minutes until approximately one week post-injury. Augmentation of the level of HO in injured vascular tissue shortly after the injury has occurred inhibits an initial increase in local VSMC proliferation post-injury. For example, such early

- 5 -

stage intervention is carried out within 24 hours post-injury.

Other features and advantages of the invention will be apparent from the following detailed description, and from the claims.

Detailed Description

The drawings will first be briefly described.

Drawings

Fig. 1 is a diagram of the targeted gene disruption strategy used in making an HO-1-deficient mouse.

Fig. 2 is a bar graph showing that hypoxia increases hematocrit in HO-1 +/+ and -/- mice.

Fig. 3 is a bar graph showing that hypoxia markedly increases ventricular weight in HO-1 -/- mice.

Fig. 4 is a diagram of the mouse model of vein graft stenosis.

Fig. 5A is a line graph showing luminal occlusion of an artery into which a vein patch has been grafted.

Fig. 5B is a diagram of a vein graft.

Fig. 6 is a diagram of plasmid containing a myosin heavy chain promoter which directs cardiospecific expression of a polypeptide-encoding DNA to which it is operably linked.

Fig. 7 is a bar graph showing that chronic hypoxia increases right ventricular systolic pressure. Wild type (+/+) and HO-1-deficient (-/-) mice were exposed to normoxia or chronic hypoxia (10% oxygen). Right ventricular pressure was measured under normoxic conditions (open bars) or after five weeks of hypoxia (filled bars). Error bars indicate standard deviation. *P<0.05 vs. animals exposed to normoxia within the same group (n = 5 in each group).

Fig. 8 is a bar graph showing that HO-1 -/- arterial smooth muscle cells are more sensitive to

- 6 -

oxidative stress compared to wild type smooth muscle cells.

Fig. 9 is an autoradiograph of a Northern blot assay showing expression of a human HO-1 (hHO-1)

5 transgene in a transgenic mouse.

Fig. 10 is an autoradiograph of a Western blot showing the presence of a hHO-1 gene product in tissues of HO-1 transgenic mice.

HO-1-deficient mice

10 HO-1-deficient (HO-1^{-/-}) mice were produced using a standard targeted gene deletion strategy to delete exon 3 (Fig. 1). The murine HO-1 gene contains 5 exons and 4 introns, spanning approximately 7 kilobases (kb). The targeting construct was made by deleting the largest exon
15 (exon 3) which contains 492 nucleotides out of the 867 nucleotides of the entire open reading frame. This deletion renders the HO-1 enzyme non-functional. An *XhoI/BamHI* fragment of the neo cassette from pMC1neo PolyA plasmid was subcloned into pBluescript II SK
20 (Stratagene, La Jolla, CA) to generate pBS-neo. To generate pBS-neo-HO-1, the 3 kb *XhoI* fragment of the HO-1 gene spanning from exon 1 to the end of intron 2 was subcloned into the *XhoI* site of pBS-neo in the same orientation as the neo cassette. The 4 kb HO-1 *BamHI-EcoRI* fragment containing a small portion of intron 3,
25 exon 4, and exon 5 was subcloned into *BamHI* and *EcoRI* site of pPGK-TK to generate pPGK-TK-HO-1. The 7 kb *BamHI-ClaI* fragment (filled in with Klenow) from pPGK-TK-HO-1 was then subcloned into *BamHI* and *XbaI* sites (filled
30 in with Klenow) sites of pBS-neo-HO-1 to generate the HO-1 targeting construct. The linearized targeting construct was transfected into murine D3 embryonic stem (ES) cells, and a clone with the correct homologous recombination (yielding the appropriately disrupted HO-1
35 gene) injected into blastocysts and used to generate HO-1

- 7 -

deficient mice. The survival rate of HO-1 $-/-$ mice was 25% of the expected survival rate, and the mice were grossly normal. The mice were deficient in HO-1 mRNA and HO-1 protein but not HO-2 mRNA or protein.

5 HO-1 transgenic mice

Standard techniques were used to generate mice which express a hHO-1 transgene in heart tissue. The transgene was cloned under the control of the cardiac α -myosin heavy chain promoter for expression preferentially
10 in cardiovascular tissue. One group of transgenic mice were engineered to express hHO-1 DNA in the sense orientation, and another group expressed hHO-1 DNA in the antisense orientation. As shown in Fig. 9, hHO-1 mRNA was detected in the heart (ventricle) of the transgenic
15 mouse but not in other tissues tested. Western blot analysis confirmed the presence of a transgenic hHO-1 gene product in heart tissue (ventricles) of hHO-1 transgenic mice. hHO-1 transgenic mice are used to evaluate the effect of HO-1 expression (and
20 overexpression) in cardiovascular tissue, e.g., in response to injury or stress.

Inhibition of cardiomyocyte death

Oxidative stress caused by such conditions as ischemia and reperfusion injury induces myocardial
25 dysfunction and cardiomyocyte death. HO is an enzyme that catalyzes oxidation of heme to generate carbon monoxide (CO; which can increase cellular cGMP) and biliverdin (which is a potent antioxidant). HO-1 is an inducible isoform of HO, whereas HO-2 is constitutively
30 expressed. Expression of HO-1 is induced in the cardiovascular system by such stimuli as hypoxia, hyperoxia, cytokines such as interleukin- 1β (IL- 1β), endotoxemia, heat shock, and ischemia. HO-1 also regulates VSMC growth. Expression of inducible HO (HO-1)

- 8 -

is markedly induced in the cardiovascular system by stimuli such as increased pressure and hypoxia.

To study the effect of HO-1 on mammalian responses to hypoxia such as that manifested in clinical conditions, e.g., high altitude pulmonary edema, myocardial infarction, myocarditis, pulmonary hypertension, pulmonary embolism, pulmonary valve stenosis, congenital heart disease, and chronic obstructive pulmonary disease (e.g., emphysema), mice were subjected to chronic hypoxia. In accordance with a standard model of pulmonary hypertension, mice were kept in a 10% O₂ chamber for 7 weeks. Two groups (Group I = five HO-1 +/+ mice; Group II = five HO-1 -/- mice) were studied. Two of the HO-1 deficient mice died at week 7; none of the HO-1 +/+ mice died. As shown in Fig. 2, exposure of the mice to hypoxic conditions resulted in an increase in hematocrit in both wild type and knockout (HO-1 -/-) mice, indicating a high level of tissue hypoxia of the mice. Under normoxia conditions, the heart weight of HO-1-deficient and wild type mice was comparable, but under hypoxic conditions the ventricular weight of HO-1-deficient mice greater than that of the HO-1-deficient mice kept under normoxic conditions (Fig. 3). For example, exposure to hypoxia for 7 weeks caused a 32% increase in the ventricular weight index in wild type mice, whereas in HO-1-deficient mice, the ventricular weight index after 7 weeks of hypoxia increased 100% (compared to normoxic wild type mice. Changes in the ventricular weight reflected mainly a right ventricular effect, as the RV(LV+septum) increased in HO-1-deficient mice compared to wild type mice exposed to hypoxia for 7 weeks.

Fig. 7 shows the effect of hypoxia on right ventricular systolic pressure, an indicator of pulmonary arterial systolic pressure. Right ventricular systolic

- 9 -

pressure in wild type and HO-1 $-/-$ mice did not differ under normoxic conditions ($P = 0.80$; Fig. 7, open bars). Although five weeks of hypoxia increased right ventricular systolic pressure, it did so to a similar
5 degree in wild type and HO-1 $-/-$ mice ($P = 0.43$; Fig. 7, filled bars).

In HO-1-deficient mice, exposure to conditions of chronic hypoxia resulted in more dramatic hypertrophy and dilation of the right ventricle of the heart compared to
10 that observed in wild type mice. Evidence of massive cardiomyocyte death was detected and large organized thrombi attached to areas of infarct were also detected in HO-1-deficient mice but not in wild type mice.

When stained with Masson's trichrome stain (which
15 detects collagen), an increase in collagen fibers was observed in tissue sections of the right ventricle of HO-1 $-/-$ mice in response to hypoxia compared to the amount of collagen detected in wild type sections. These data indicate evidence of scar formation and repair mechanisms
20 in the hearts of HO-1-deficient mice under hypoxic conditions. Tissue sections were also stained with PECAM stain (which detects endothelial cells). Increased blood vessel formation was detected in the right ventricles of HO-1 $-/-$ mice in response to hypoxia
25 compared to wild type mice. These data suggest that HO-1 inhibits angiogenesis in response to hypoxia.

The mechanism of cardiomyocyte death in HO-1 $-/-$ mice under hypoxic conditions was evaluated by histological analysis, immunocytochemistry, and TdT-mediated dUTP-biotin nickend labeling (TUNEL assay). The
30 standard TUNEL assay detects apoptosis. Ventricles were fixed in 4% paraformaldehyde overnight at 4°C and embedded in paraffin. Tissue sections were stained with hematoxylin and eosin or Masson's trichrome. To detect
35 oxidation-specific lipid-protein adducts, heart tissue

- 10 -

sections were immunostained with polyclonal antibody MAL-2 (anti-malondialdehyde-lysine; Rosenfeld et al., 1990, Arteriosclerosis 10:336-349) and counterstained with methyl green. TUNEL was used to detect DNA breaks in apoptotic cells *in situ*. Red staining in the nuclei of the cells indicated a positive reaction; the cells were also counterstained with methyl green. Pulmonary vascular remodeling was assessed in lungs that had been perfused with saline through the pulmonary artery and fixed with 4% paraformaldehyde instilled through the trachea. Muscularization of peripheral vessels was determined using standard methods, e.g., that described in Klinger et al., 1993, J. Appl. Physiol. 75:198-205. The nuclei of cardiomyocytes of HO-1-deficient mice subjected to hypoxic conditions stained positive, providing evidence that apoptosis contributes to the mechanism of death of cardiomyocytes under these conditions.

Additional histological analyses were undertaken to confirm that chronic hypoxia induces right ventricular infarction in HO-1-deficient mice. Cardiomyocytes were intact in ventricular sections from wild type mice exposed to 7 weeks of hypoxia, but ventricular sections from HO-1-deficient mice exposed to 7 weeks of hypoxia showed mononuclear inflammatory cell infiltration, extensive cardiomyocyte degeneration, and death with focal calcification. These observations indicate that infarcts were 1-2 weeks old. The right ventricular infarcts did not appear to result from vascular occlusion, because the coronary arteries supplying blood to the right ventricle were patent in HO-1-deficient mice.

To detect collagen accumulation indicative of fibrosis, ventricular sections were stained with Masson's trichrome. After 7 weeks of hypoxia, cells surrounding

- 11 -

blood vessels stained positive for collagen in hearts from wild type mice. In hearts from HO-1-deficient mice exposed to hypoxia for 7 weeks, early collagen deposition was present throughout the lesion. The degree of
5 fibrosis was consistent with early scar formation and a lesion of 1-2 weeks old.

Two out of the six hearts from 7-week hypoxic HO-1-deficient mice did not exhibit right ventricular infarcts; however, close examination of the hearts
10 revealed focal areas of myocardial degeneration without evidence of extensive inflammatory cell infiltration. These hearts showed early evidence of myocardial damage. Heart from wild type and HO-1-deficient mice examined after 5 weeks of hypoxia displayed no cardiomyocyte
15 degeneration or death and no extensive mononuclear inflammatory cell infiltration. Cells surrounding blood vessels stained positive for collagen in hearts from wild type and HO-1-deficient mice housed for 5 weeks under normoxic and hypoxic conditions. In contrast with the
20 right ventricular free walls from HO-1-deficient mice exposed to 7 weeks of hypoxia (which showed deposition of collagen), no collagen deposition was evident in the right ventricular free walls from HO-1-deficient mice exposed to 5 weeks of hypoxia. These data confirm that
25 in the HO-1-deficient mice exposed to 7 weeks of hypoxia, myocardial infarcts were less than 2 weeks old.

To assess oxidative damage in hearts with infarcts taken from HO-1-deficient mice, ventricular tissue sections were immunostained with MAL-2 which detects
30 oxidation-specific lipid-protein adducts. In contrast to the minimal MAL-2 staining observed in hearts from wild type mice, MAL-2 staining was intense in cells (predominantly cardiomyocytes) beneath the infarcted area of the right ventricle in HO-1-deficient mice. These
35 data indicate the presence of severe oxidative damage

- 12 -

within and around the infarct site. No TUNEL-positive cardiomyocytes were detectable in right ventricles from wild-type mice, but a significant number of TUNEL-positive cells surrounded the infarct site in right
5 ventricles form HO-1-deficient mice.

The data described herein indicate that

(1) pulmonary vascular remodeling in response to hypoxia is similar in HO-1 +/+ and -/- mice, (2) hypoxia induces more severe right ventricular hypertrophy in HO-1 -/-mice
10 than in HO+/+ mice, and (3) in HO-1 -/- (but not +/+ mice), massive cardiomyocyte death occurs with large organized thrombi attached to the infarct site. Although some cardiomyocyte death appears to be due to necrosis, apoptosis is a significant mechanism of cardiomyocyte
15 death. Hypoxia and elevated pulmonary arterial pressure increase cardiac production of reactive oxygen species, which play a significant role in myocardial death during ischemia/reperfusion.

Although myocardial infarction has been shown to
20 increase oxidative stress, a 2-3 fold increase in the nitration of protein tyrosine residues (which indicates the presence of the potent oxidant peroxynitrite) was detected in noninfarcted HO-1-deficient hearts exposed to 7 weeks of hypoxia. These data indicate that an increase
25 in oxidative stress precedes gross myocardial infarction.

Evidence of extensive lipid peroxidation in the zone of right ventricular infarction supports the conclusion that the absence of HO-1 in cardiomyocytes leads to an accumulation of reactive oxygen species that
30 causes cardiomyocyte death. Administration of HO-1 or overexpression of HO-1 protects against cardiomyocyte damage from hemodynamic stress or ischemia/reperfusion.

Adaptation of the cardiovascular system to hypoxia

The gene deletion studies described herein
35 indicate that HO-1 plays an important protective role in

- 13 -

vivo in the adaptation of the cardiovascular system to hypoxia. Right ventricles from HO-1 -/- mice were severely dilated and contained right ventricular infarcts with mural thrombi.

- 5 Humans and animals respond to hypoxia by exhibiting pulmonary vascular remodeling, pulmonary hypertension, and hypertrophy of the right ventricle. The data described herein were obtained using a mouse model of vascular injury which mimics the human response.
- 10 Hypoxia induces HO-1 expression in the lung, and CO generated by hypoxic VSMCs inhibits proliferation of these cells.

The data described herein indicate that the absence of HO-1 results in a maladaptive response in

15 cardiomyocytes exposed to hypoxia-induced pulmonary hypertension. In the absence of HO-1, VSMC are more sensitive to oxidative stress and have a maladaptive response to pressure overload. HO-1 has a protective effect on cardiomyocytes and VSMC subjected to stress

20 such as pressure-induced injury and secondary oxidative damage.

Therapeutic administration of HO

- In the absence of HO-1, cardiomyocytes undergo apoptotic cell death when subjected to stress such as
- 25 pressure overload or exposure to reactive oxygen species and that death can be inhibited by contacting the cells with HO. For example, HO-1, HO-2, or HO-3 protein or polypeptide (or DNA encoding HO-1, HO-2, or HO-3) is administered locally to heart tissue affected by hypoxic
- 30 conditions. One means for accomplishing local delivery is providing an HO or DNA encoding and HO on a surface of a vascular catheter, e.g., a balloon catheter coated with an antioxidant, which contacts the wall of the blood vessel to deliver therapeutic compositions at the site of

- 14 -

contact. Drug delivery catheters can also be used to administer solutions of therapeutic compositions.

HO-1 is therapeutically overexpressed (e.g., by administering an inducing agent to increase expression
5 from the endogenous gene) or by administering DNA (alone or in a plasmid) encoding an HO such as HO-1 or HO-2 (or an active fragment thereof, i.e., a fragment has the activity of inhibiting cardiomyocyte death). Inducing agents that stimulate HO-1 expression in cells include
10 hemin, hemoglobin, and heavy metals, e.g., SnCl_2 or NiCl_2 . For example, 250 mmol/kg of body weight of SnCl_2 or NiCl_2 is administered subcutaneously or 15 mg/kg of body weight of hemin is administered intraperitoneally to laboratory animals. Doses for human patients are determined and
15 optimized using standard methods.

Tables 1 and 3 show human HO-1 and HO-2 cDNA, respectively, in which the polypeptide-encoding nucleotides are designated in bold type and the termination codon is underlined. Tables 2 and 4 show the
20 amino acid sequences of human HO-1 and HO-2, respectively. Tables 5 and 6 show the nucleotide and amino acid sequence of rat HO-3.

- 15 -

TABLE 1: Human HO-1 cDNA

1 tcaacgcctg cctcccctcg agcgtcctca ggcagaccgc
cgccccgcga gccagcacga
61 acgagcccag caccggccgg atggagcgtc cgcaaccgga
5 cagcatgccc caggatttgt
121 cagaggccct gaaggaggcc accaaggagg tgcacaccca
ggcagagaat gctgagttca
181 tgaggaactt tcagaagggc caggtgaccc gagacggctt
caagctgggtg atggcctccc
10 241 tgtaccacat ctatgtggcc ctggaggagg agattgagcg
caacaaggag agcccagtct
301 tcgcccctgt ctacttccca gaagagctgc accgcaaggc
tgccctggag caggacctgg
361 ccttctggta cgggccccgc tggcaggagg tcatccccta
15 cacaccagcc atgcagcgt
421 atgtgaagcg gctccacgag gtggggcgca cagagcccga
gctgctgggtg gcccacgcct
481 acaccgcta cctgggtgac ctgtctgggg gccaggtgct
caaaaagatt gcccagaaag
20 541 ccctggacct gcccagctct ggcgagggcc tggccttctt
caccttcccc aacattgcca
601 gtgccaccaa gttcaagcag ctctaccgct cccgcatgaa
ctccctggag atgactcccg
661 cagtcaggca gaggggtgata gaagaggcca agactgcgtt
25 cctgctcaac atccagctct
721 ttgaggagtt gcaggagctg ctgacccatg acaccaagga
ccagagcccc tcacgggcac
781 cagggtctcg ccagcgggcc agcaacaaag tgcaagattc
tgcccccggtg gagactccca
30 841 gaggggaagcc cccactcaac acccgctccc aggctccgct
tctccgatgg gtccttacac
901 tcagctttct ggtggcgaca gttgctgtag ggctttatgc
catgtgaatg caggcatgct

- 16 -

961 ggctcccagg gccatgaact ttgtccggtg gaaggccttc
tttctagaga gggaattctc
1021 ttggetggct tccttaccgt gggcactgaa ggctttcagg
gcctccagcc ctctcactgt
5 1081 gtccctctct ctggaaagga ggaaggagcc tatggcatct
tccccaacga aaagcacatc
1141 caggcaatgg cctaaacttc agagggggcg aaggggtcag
ccctgccctt cagcatcctc
1201 agttcctgca gcagagcctg gaagacaccc taatgtggca
10 gctgtctcaa acctccaaaa
1261 gccctgagtt tcaagtatcc ttgttgacac ggccatgacc
actttccccg tgggccatgg
1321 caatttttac acaaacctga aaagatgttg tgtcttgtgt
ttttgtctta tttttgttgg
15 1381 agccactctg ttcttggtc agcctcaa atgcagtatttt
tggtgtgttc tggtgtttt
1441 atagcagggt tgggggtggt tttgagccat gcgtgggtgg
ggagggaggt gtttaacggc
1501 actgtggcct tgggtctaact tttgtgtgaa ataataaaca
20 acattgtctg
(SEQ ID NO:1)

Table 2: Human HO-1 amino acid sequence

MERPQPDSMP QDLSEALKEA TKEVHTQAEN AEFMRNFQKG QVTRDGFKL
MASLYHIYVA
25 LEEEIERNKE SPVFAPVYFP EELHRKAALE QDLAFWYGPR WQEVIPYTPA
MORYVKRLHE
VGRTEPELLV AHAYTRYLGD LSGGQVLKKI AQKALDLPSS GEGLAFFTFP
NIASATKFKQ
LYRSRMNSLE MTPAVRQRVI EEAKTAFLN IQLFEELQEL LTHDTKDQSP
30 SRAPGLRQRA
SNKVQDSAPV ETPRGKPPLN TRSQAPLLRW VLTLNFLVAT VAVGLYAM (SEQ
ID NO:2)

- 17 -

Table 3: Human HO-2 cDNA

1 gggctgactg gaggctggcg gacaggcgac agacctgagg
caggaccaga ggagcgagac
61 gagcaagaac cacacccagc agcaatgtca gcggaagtgg
5 aaacctcaga gggggtagac
121 gagtcagaaa aaaagaactc tggggcccta gaaaaggaga
accaaattgag aatggctgac
181 ctctcagagc tcctgaagga agggaccaag gaagcacacg
accgggcaga aaacacccag
10 241 tttgtcaagg acttcttgaa aggcaacatt aagaaggagc
tgtttaagct ggccaccacg
301 gcactttact tcacatactc agccctcgag gaggaaatgg
agcgcaacaa ggaccatcca
361 gcctttgccc ctttgtactt ccccatggag ctgcaccgga
15 aggaggcgct gaccaaggac
421 atggagtatt tctttggtga aaactgggag gagcaggtgc
agtgccecaa ggctgcccag
481 aagtacgtgg agcggatcca ctacataggg cagaacgagc
cggagctact ggtggcccat
20 541 gcatacacc gctacatggg ggatctctcg gggggccagg
tgctgaagaa ggtggcccag
601 cgagcactga aactccccag cacaggggaa gggaccagt
tctacctgtt tgagaatgtg
661 gacaatgccc agcagttcaa gcagctctac cgggccagga
25 tgaacgccct ggacctgaac
721 atgaagacca aagagaggat cgtggaggcc aacaaggctt
ttgagtataa catgcagata
781 ttcaatgaac tggaccaggc cggctccaca ctggccagag
agaccttgga ggatgggttc
30 841 cctgtacacg atgggaaagg agacatgcgt aatgccctt
tctacgctgc tgaacaagac
901 aaagggtcgg agggcagcct gtcccttcg acaagctatg
ctgtgctgag gaagcccagc

- 18 -

961 ctccagttca tcttgccgc tgggtgtggcc ctagctgctg
gactcttggc ctggtactac
1021 atgtgaagca cccatcatgc cacaccgga cctcctccc
gactgaccac tggcctaccc
5 1081 ctttctccag cctgactaa actaccacct caggtgactt
tttaaaaaat gctgggttta
1141 agaaaggcaa ccaataaaag agatgctaga gcctcgtctg
acagcatcct ctctatgggc
1201 catattccgc actgggcaca ggccgtcacc ctgggagcag
10 tgggcacagt gcagcaagcc
1261 tggccccga ccagctcta ctccaggctt ccacacttct
gggccctagg ctgcttcgg
1321 tagtcctgt ttttgcagta catgggtgac tatctccct
gttggagggtg agtggcctgt
15 1381 aagtccaagc tgtgcgaggg ggccttgctg gatgctgctg
tacaacttct ggcctctct
1441 tggaccctgg gagtgagggt ggggtgtgggt ggaagcctca
gaggccttgg gagctcatcc
1501 ctctcaccca gaatccctct aacccttggg tgcggtttgc
20 tcagccccag cttatctcct
1561 cctccgctg tgtaaagtct ccagcactca ataaagtggg
ctttgcaagc taataaaaaa
1621 aaaaaaa (SEQ ID NO:3)

- 19 -

Table 4: Human HO-2 amino acid sequence

MSAEVETSEGVDESEKKNSGALEKENQMRMADLSELLKEGTKEAHDRAENTQFVKDF
LKGNIKKELFKLATTALYFTYSALEEEMERNKDHFAFAPLYFPMELHRKEALTKDME
YFFGENWEEQVQCPKAAQKYVERIHYYIGQNEPELLVAHAYTRYMGDLSGGQVLKKVA
5 QRALKLPSTGEGTQFYLFENVDNAQQFKQLYRARMNALDLNMKTKERIVEANKAFEY
NMQIFNELDQAGSTLARETLEDGFPVHDGKGDMRKCPFYAAEQDKGLEGSLSLPTSY
AVLRKPSLQFILAAGVALAAGLLAWYYM (SEQ ID NO:4)

Table 5: Rat HO-3 nucleotide sequence

1 tttcagggat ttttgcgatt cctctctgta gacttctact
10 tgttctctaa gggagttctt
61 catgtctttc ttgaagtcac ccagcatcat gatcaaatat
gattttgaaa ctagatcttg
121 cttttctggt gtgtttggat attccatggt tgttttggtg
ggagaattgg gctccgatga
15 181 tggcatgtag tcttggtttc tgttgcttgg tttcctgcgc
ttgcctctcg ccatcagatt
241 atctctagtg ttactttggt ctgctatttc tgacagtggc
tagactgtcc tataagcctg
301 tgtgtcagga gtgctgtaga ccttttttcc tctctttcag
20 tcagttatgg gacagagtgt
361 tctgcttttg ggcgtgtagt ttttctctc tacaggtctt
cagctgttcc tgtgggcctg
421 tgtcttgagt tcaccaggca gctttcttgc agcagaaaat
ttggtcatac ctgtgatcct
25 481 gaggctcaag ttcgctcgtg ggggtgctgc caggggctct
ctgcagcggg cacaaccagg
541 aagacctgtg cggccccttc cggagcttca gtgcaccagg
gttccagatg gcctttggcg
601 ttttctctg gcgtccgaga tgtatgtaca gagagcagtc
30 tcttctgggt tcccaggctt
661 gtctgctct ctgaagggtc agctctccct cccacgggat
ttgggtgcag agaactgtt

- 20 -

721 atccggtctg tttctttcag gttccggtgg tgtctcaggc
aggtgtcgtt cctgcgcct

781 ccccatggg accagaggcc ttatacagtt tcctcttggg
ccagggatgt gggcaggggt

5 841 gagcagtgtt ggtggtctct tccgtctgca gcctcaggag
tgccacctga ccaggcgggt

901 gggctctctct ctgagaattt cattttttaa tcattcatta
aatgtcatg acttgatgtc

961 ctgctgtccg tctcacgcc tcagctgtaa cagtgccgag
10 ggagtactg aagaagagac

1021 tgaatgacca gagtatgggc agcacagaca actcaacaaa
aatgtcttca gaggtggaga

1081 ctgcggaggc cgtagatgag tcagagaaga actctatggc
atcagagaag gaaaaccatt

15 1141 ccaaatagc agacttttct gatcttctga aggaaggagc
aaaggaagca gatgaccggg

1201 cagaaaatac ccagtttgtc aaagacttct tgaaaggaaa
cattaagaag gagctattta

1261 agctggccac cactgcactt tcatactcag cccctgagga
20 ggaaatggat tcaactgacca

1321 aggacatgga gtacttcttt ggtgaaaact gggaggaaaa
agtgaagtgc tctgaagctg

1381 cccagacgta tgtggatcag attcactatg tagggcaaaa
tgagccagag catctgggtg

25 1441 cccatactta ctctacttac atgggggggaa acctttcagg
ggaccaggta ctgaagaagg

1501 agaccagcc ggtccccttc actagggaag ggactcagtt
ctacctgttt gagcatgtag

1561 acaatgctaa gcaattcaag ctattctact gcgctagatt
30 gaatgccttg gacctgaatt

1621 tgaagaccaa agagaggatt gtggaggaag ccaccaaacg
ctttgaatat aatatgcaga

1681 tattcagtga actggaccag gcaggctcca taccagtaag
agaaacccta aagaatgggc

- 21 -

1741 tctcaatact tgatgggaag ggaggtgtat gcaaagtgtcc
 ctttaatgct gctcagccag
 1801 acaaaggtac cctgggaggc agcaactgcc ctttccagat
 gtccatggcc ttgctgagga
 5 1861 agcctaactt gcagctcatt ctagttgcca gtatggcctt
 ggtagctgga cttttagcct
 1921 ggtactacat gtgaagggcc tgtcaagttg tttgcatcct
 atctcaacat cctaccactt
 1981 gttccttccc cacctccacc tctgcctaga actaccacct
 10 caggtgacat ttttaatgtt
 2041 gggtttgaga aaatgagcaa ccaataaaaag acagacccta
 gaaaaaagtc atgacttaag
 2101 tggcacgggg acacctaaag tcacactttg tgcttcagac
 atactttctt tctctatttc
 15 2161 aacactgaat tcgggaagta acctactact attaataata
 aatgctacac aatgcataat
 2221 aaaaa (SEQ ID NO:5)

Table 6: Rat HO-3 amino acid sequence

MSSEVETAEAVDESEKNSMASEKENHSKIADFSDLLKEGTKEADDRAENTQFVKDFL
 20 KGNIKKELFKLATTALSYSAPEEEMDSLTKDMEYFFGENWEEKVKCSEAAQTYVDQI
 HYVGQNEPEHLVAHTYSTYMGGNLSGDQVLKKETQVPVFTREGTQFYLFVHDNAKQ
 FKLPHYCARLNALDLNLKTKERIVEEATKAFFEYNMQIFSELDQAGSIPVRETLKNGLS
 ILDGKGGVCKCPFNAAPDKGTLGGSNCPFQMSMALLRKPNLQLILVASMALVAGLL
 AWYYM (SEQ ID NO:6)

25 An HO preferably has an amino acid sequence that
 is at least 85% identical (preferably at least 90%, more
 preferably at least 95%, more preferably at least 98%,
 most preferably at least 100% identical) to the amino
 acid sequence of SEQ ID NO: 2, 4, or 6. DNA encoding an
 30 HO preferably has nucleotide sequence that is at least
 50% identical (preferably at least 75%, more preferably
 at least 85%, more preferably at least 95%, most

- 22 -

preferably at least 100% identical) to the nucleotide sequence of the coding region of SEQ ID NO:1, 3, or 5.

The per cent identity of nucleotide and amino acid sequences is determined using the Sequence Analysis

5 Software Package developed by the Genetics Computer Group (University of Wisconsin Biotechnology Center, Madison, WI), employing the default parameters thereof.

To be clinically beneficial, expression of HO from an endogenous gene or expression of recombinant HO from
10 exogenous DNA need not be long term. For example, to inhibit cardiomyocyte death associated with a myocardial infarction or other acute condition which results in oxidative stress to the tissue, the most critical period of treatment is the first three months after injury.

15 Thus, gene therapy to express recombinant HO for even a period of days or weeks after administration of the HO-encoding DNA (which administration is prior to or soon after an injury) minimizes cell death, inhibits VSMC proliferation, and therefore confers a clinical benefit.

20 For local administration of DNA to cardiovascular tissue, standard gene therapy vectors used. Such vectors include viral vectors, including those derived from replication-defective hepatitis viruses (e.g., HBV and HCV), retroviruses (see, e.g., WO 89/07136; Rosenberg et
25 al., 1990, N. Eng. J. Med. 323(9):570-578), adenovirus (see, e.g., Morsey et al., 1993, J. Cell. Biochem., Supp. 17E), adeno-associated virus (Kotin et al., 1990, Proc. Natl. Acad. Sci. USA 87:2211-2215), replication
defective herpes simplex viruses (HSV; Lu et al., 1992,
30 Abstract, page 66, Abstracts of the Meeting on Gene Therapy, Sept. 22-26, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York), and any modified versions of these vectors. The invention may utilize any other delivery system which accomplishes *in vivo* transfer of
35 nucleic acids into eukaryotic cells. For example, the

- 23 -

nucleic acids may be packaged into liposomes, e.g., cationic liposomes (Lipofectin), receptor-mediated delivery systems, non-viral nucleic acid-based vectors, erythrocyte ghosts, or microspheres (e.g.,
5 microparticles; see, e.g., U.S. Patent No. 4,789,734; U.S. Patent No. 4,925,673; U.S. Patent No. 3,625,214; Gregoriadis, 1979, *Drug Carriers in Biology and Medicine*, pp. 287-341 (Academic Press,)). Naked DNA may also be administered. Alternatively, a plasmid which directs
10 cardiospecific expression (e.g., a plasmid containing a myosin heavy chain (α MHC) promoter; Fig. 6) of an HO-encoding sequence can be used for gene therapy. Expression of an HO (encoded, e.g., by the coding sequences of SEQ ID NO:1, 3, or 5) from such a
15 constitutive promoter is useful to inhibit cardiomyocyte death *in vivo*. Nucleic acids which hybridize at high stringency to the coding sequences of SEQ ID NO:1, 3, or 5 and which encode a polypeptide which has a biological activity of an HO polypeptide (e.g., inhibition of
20 cardiomyocyte death) are also used for gene therapy for vascular injury. To determine whether a nucleic acid hybridizes to a reference nucleic acid at a given stringency, hybridization is carried out using standard techniques, such as those described in Ausubel et al.
25 (*Current Protocols in Molecular Biology*, John Wiley & Sons, 1989). "High stringency" refers to nucleic acid hybridization and wash conditions characterized by high temperature and low salt concentration, e.g., wash conditions of 65°C at a salt concentration of
30 approximately 0.1 X SSC. "Low" to "moderate" stringency refers to DNA hybridization and wash conditions characterized by low temperature and high salt concentration, e.g., wash conditions of less than 60°C at a salt concentration of at least 1.0 X SSC. For example,
35 high stringency conditions may include hybridization at

- 24 -

about 42°C, and about 50% formamide; a first wash at about 65°C, about 2X SSC, and 1% SDS; followed by a second wash at about 65°C and about 0.1% x SSC. Lower stringency conditions suitable for detecting DNA

5 sequences having about 50% sequence identity to an CHF-1 gene are detected by, for example, hybridization at about 42°C in the absence of formamide; a first wash at about 42°C, about 6X SSC, and about 1% SDS; and a second wash at about 50°C, about 6X SSC, and about 1% SDS. To
10 determine whether a polypeptide encoded by a hybridizing nucleic acid has a biological activity of an HO polypeptide, the polypeptide is evaluated using any of the functional assays to measure HO activity described herein, e.g., measuring VSMC proliferation or
15 cardiomyocyte death.

For gene therapy of cardiovascular tissue, fusigenic viral liposome delivery systems known in the art (e.g., hemagglutinating virus of Japan (HVJ) liposomes or Sendai virus-liposomes) are useful for
20 efficiency of plasmid DNA transfer (Dzau et al., 1996, Proc. Natl. Acad. Sci. U.S.A. 93:11421-11425). Using HVJ-liposomes, genes are expressed from plasmid DNA delivered to target tissues in vivo for extended periods of time (e.g., greater than two weeks for heart and
25 arterial tissue and up to several months in other tissues).

DNA for gene therapy can be administered to patients parenterally, e.g., intravenously, subcutaneously, intramuscularly, and intraperitoneally.
30 Sustained release administration such as depot injections or erodible implants, e.g., vascular stents coated with DNA encoding an HO, may also be used. The compounds may also be directly applied during surgery, e.g., bypass surgery, or during angioplasty, e.g., an angioplasty
35 catheter may be coated with DNA encoding an HO. The DNA

- 25 -

is then deposited at the site of angioplasty. DNA or an inducing agent is administered in a pharmaceutically acceptable carrier, i.e., a biologically compatible vehicle which is suitable for administration to an animal e.g., physiological saline. A therapeutically effective amount is an amount which is capable of producing a medically desirable result, e.g., expression of HO, in a treated animal. Such an amount can be determined by one of ordinary skill in the art. As is well known in the medical arts, dosage for any given patient depends upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, severity of arteriosclerosis or vascular injury, and other drugs being administered concurrently. Dosages will vary, but a preferred dosage for intravenous administration of DNA is approximately 10^6 to 10^{22} copies of the DNA molecule.

HO-based therapy for cardiovascular disorders depends on when (in the course of a vascular injury) the patient is encountered. HO-1 -/- VSMC initially proliferated at a faster rate compared to wild type VSMC within days after an insult. Increasing local HO-1 levels at this stage inhibits VSMC growth and confers a clinical benefit. For example, if the patient is encountered at an early stage (e.g., within one week of cardiovascular stress or injury), the patient is treated by augmenting the local level of HO-1 (e.g., by administering an HO polypeptide, by increasing expression of endogenous HO, or by standard gene therapy techniques described above to produce recombinant HO *in vivo*) to inhibit the growth of VSMC and decrease the size of a myocardial infarct. In contrast, if the patient is encountered at a later stage (e.g., several weeks, 1 month, 2 months, and up to 3 months after an injury), the

- 26 -

patient is treated by inhibiting HO expression to decrease formation of a stenotic lesion, e.g., by antisense therapy, as described below.

Organ and tissue preservation

5 Concerns about irreversible ischemic tissue damage arise when a donor organ, e.g., a heart, is removed from the donor and stored for more than about 5-6 hours before transplantation into the recipient. *Ex vivo* treatment of a donor organ to reduce tissue damage by inhibiting death
10 of cardiomyocytes is carried out by immersing the organ in a solution containing an inducing agent, an HO, e.g., HO-1 or HO-2, or a nucleic acid encoding an HO prior to transplantation. By "ex vivo treatment" is meant treatment that takes place outside of the body. For
15 example, ex vivo treatment is administered to an organ (or a tissue fragment or dissociated cells) that has been removed from a mammal and which will be returned to the same or a different mammal (at the same or different anatomical site) after the treatment. For example,
20 effective DNA delivery to cells in solid organs achieved by contacting the organ with a combination of DNA, liposomes and transferrin during the cold ischemic time prior to transplantation (see, e.g., Hein et al. 1998, Eur. J. Cardiothorac. Surg. 13:460-466). An organ may
25 also be perfused or electroporated with solution containing HO-encoding DNA. Vectors and delivery systems described above for gene therapy applications are suitable for organ preservation and cell preservation *in vitro*.

30 Inhibition of restenosis

VSMC proliferation contributes to graft stenosis and restenosis following vascular injury such as that resulting from coronary angioplasty and coronary bypass surgery. Patients with restenosis have a significantly

- 27 -

poorer clinical outcome compared to patients without restenosis.

Using a mouse model of vascular graft stenosis in which the stenosis develops rapidly and closely mimics the development of vascular graft stenosis in humans, the effect of HO-1 on VSMC proliferation was examined. A patch of jugular vein was grafted onto a carotid artery in normal and HO-1 deficient mice to create composite vessels that mimic vein grafts used for bypass surgery (Fig. 4). The vein patch is subject to increased pressure which leads to an increase in local VSMC proliferation and occlusion of the blood vessel (Figs. 5A-B). In wild type mice, there was robust formation of a neointima (characterized by proliferating VSMC) in the vein graft. In contrast, tissue sections of the neointima of HO-1 -/- mice revealed a necrotic mass. The HO-1 -/- neointima was a complex lesion characterized by mostly acellular material, indicating death of VSMC. HO-1 -/- VSMC are more susceptible to H₂O₂-induced death compared to VSMC isolated from wild type mice (Fig. 8). These data indicate that HO-1 is required for VSMC proliferation at later stages post-injury and that local inhibition of HO-1 expression, e.g., by antisense therapy, is useful to inhibit graft stenosis or restenosis in patients affected or who at risk of developing such conditions.

The data described herein indicate that (1) in response to increased pressure, VSMC proliferate in the neointima of the venous patch in HO-1 +/+ mice, and (2) in contrast, massive cell death occurs in the neointima of the venous patch in HO-1 -/- mice.

Patients undergoing invasive vascular procedures, e.g., balloon angioplasty, are at risk for developing undesired vascular stenosis or restenosis. Angioplasty, used to treat arteriosclerosis, involves the insertion of

- 28 -

catheters, e.g., balloon catheters, through an occluded region of a blood vessel in order to expand it. However, the aftermath of angioplasty may be problematic.

Restenosis, or closing of the vessel, can occur as a

5 consequence of injury, e.g., mechanical abrasion associated with the angioplasty treatment. This restenosis is caused by proliferation of smooth muscle cells stimulated by vascular injury. Other anatomical disruptions or mechanical disturbances of a blood
10 vessel, e.g., laser angioplasty, coronary artery surgery, atherectomy, coronary artery stents, and coronary bypass surgery, may also cause vascular injury and subsequent proliferation of smooth muscle cells.

Therapeutic approaches, such as antisense therapy
15 or ribozyme therapy are used to inhibit HO expression, and as a result, VSMC proliferation that leads to neointimal thickening. The antisense strand (either RNA or DNA) is directly introduced into the cells in a form that is capable of binding to the mRNA transcripts.
20 Alternatively, a vector-containing sequence which, which once within the target cells is transcribed into the appropriate antisense mRNA, may be administered. Nucleic acids complementary to all or part of the HO cDNA (SEQ ID NO: 1, 3, or 5) may be used in methods of antisense
25 treatment to inhibit expression of HO. Antisense treatment is carried out by administering to a mammal, such as a human, DNA containing a promoter, e.g., a cardiospecific promoter, operably linked to a DNA sequence (an antisense template), which is transcribed
30 into an antisense RNA. By "operably linked" is meant that a coding sequence and a regulatory sequence(s) (i.e., a promoter) are connected in such a way as to permit gene expression when the appropriate molecules (e.g., transcriptional activator proteins) are bound to
35 the regulatory sequence(s). Alternatively, as mentioned

- 29 -

above, antisense oligonucleotides may be introduced directly into vascular cells. The antisense oligonucleotide may be a short nucleotide sequence (generally at least 10, preferably at least 14, more preferably at least 20 (e.g., at least 30), and up to 100 or more nucleotides) formulated to be complementary to a portion, e.g., the coding sequence, or all of HO mRNA. Oligonucleotides complementary to various portions of HO-1 or HO-2 mRNA can readily be tested *in vitro* for their ability to decrease production of HO in cells, using standard methods. Sequences which decrease production of HO message in *in vitro* cell-based or cell-free assays can then be tested *in vivo* in rats or mice to determine whether HO expression (or VSMC proliferation) is decreased.

Ribozyme therapy can also be used to inhibit gene expression. Ribozymes bind to specific mRNA and then cut it at a predetermined cleavage point, thereby destroying the transcript. These RNA molecules may be used to inhibit expression of a gene encoding a protein involved in the formation of vein graft stenosis according to methods known in the art (Sullivan et al., 1994, J. Invest. Derm. 103:85S-89S; Czubayko et al., 1994, J. Biol. Chem. 269:21358-21363; Mahieu et al, 1994, Blood 84:3758-65; Kobayashi et al. 1994, Cancer Res. 54:1271-1275).

Standard methods of administering antisense therapy have been described (see, e.g., Melani et al., 1991, Cancer Res. 51:2897-2901). Antisense nucleic acids which hybridize to HO-encoding mRNA can decrease or inhibit production of HO by associating with the normally single-stranded mRNA transcript, thereby interfering with translation and thus, expression of HO. Such nucleic acids are introduced into target cells by standard vectors and/or gene delivery systems such as those

- 30 -

described above for gene therapy. Suitable gene delivery systems may include liposomes, receptor-mediated delivery systems, naked DNA, and viral vectors such as herpes viruses, retroviruses, adenoviruses and adeno-associated viruses, among others. For example, antisense oligodesoxynucleotides, e.g., oligonucleotides which have been modified to phosphorthioates or phosphoamidates, withstand degradation after delivery and have been successfully used to inhibit gene expression in a model of reperfusion injury (see, e.g., Haller et al., 1998, Kidney Int. 53:1550-1558). Pharmaceutically acceptable carriers are biologically compatible vehicles which are suitable for administration to an animal: e.g., physiological saline. A therapeutically effective amount of a compound is an amount which is capable of producing a medically desirable result in a treated animal, e.g., inhibition of expression of HO-1 or a decrease in VSMC proliferation.

Compositions that inhibit HO activity, e.g., its role in the promotion of VSMC proliferation, are also administered to inhibit VSMC-mediated stenosis or restenosis. For example, metalloporphyrins, e.g., zinc protoporphyrin IX (ZnPP), zinc mesoporphyrin IX (ZnMP), tin protoporphyrin IX (SnPP), tin mesoporphyrin IX (SnMP), zinc deuteroporphyrin IX 2,4 bis glycol (ZnDPBG), chromium protoporphyrin (CrPP), cobalt protoporphyrin (CoPP), and manganese metalloporphyrin (MnPP) are administered to mammals at $\mu\text{mol/kg}$ doses to inhibit HO activity. SnPP has safely been administered to human infants at doses of $0.5 \mu\text{mol/kg}$ to $100 \mu\text{mol/kg}$ of body weight. HO-inhibitory doses for local administration are determined using methods known in the art.

Parenteral administration, such as intravenous, subcutaneous, intramuscular, and intraperitoneal delivery routes, may be used to deliver the compounds that inhibit

- 31 -

HO activity or expression, with local vascular administration being the preferred route. Dosages for any one patient depends upon many factors, including the patient's size, body surface area, age, the particular
5 compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently. For antisense therapy, a preferred dosage for administration of nucleic acids is from approximately 10^6 to 10^{22} copies of the nucleic acid
10 molecule. As described above, local administration to a site of vascular injury or to cardiac tissue is accomplished using a catheter or indwelling vascular stent.

Other embodiments are within the following claims.
15 What is claimed is:

- 32 -

1. A method of inhibiting cardiomyocyte death in a mammal comprising locally administering to the myocardium of said mammal a heme oxygenase (HO).

2. The method of claim 1, wherein said mammal has suffered a myocardial infarction.

3. The method of claim 1, wherein said mammal has myocarditis.

4. The method of claim 1, wherein said HO is heme oxygenase-1 (HO-1).

5. The method of claim 1, wherein said HO is heme oxygenase-2 (HO-2).

6. A method of inhibiting cardiomyocyte death in a mammal comprising locally administering to the myocardium of said mammal a DNA encoding a HO.

7. The method of claim 6, wherein said HO is HO-1.

8. The method of claim 6, wherein said HO is HO-2 or HO-3.

9. A method of inhibiting cardiomyocyte death in vitro, comprising contacting cardiomyocytes with an HO.

10. A method of inhibiting cardiomyocyte death in vitro, comprising contacting cardiomyocytes with DNA encoding an HO.

- 33 -

11. The method of claim 10, wherein said HO is HO-1.

12. The method of claim 10, wherein said HO is HO-2.

5 13. A method of preserving isolated myocardial tissue comprising perfusing said tissue with a solution comprising an HO or a DNA encoding an HO.

10 14. A method of inhibiting vascular restenosis in a mammal comprising locally administering to the site of a vascular injury a compound which inhibits expression of HO-1.

15 15. The method of claim 14, wherein said compound inhibits HO-1 transcription in a vascular cell of said mammal.

16 16. The method of claim 15, wherein said vascular cell is an aortic smooth muscle cell.

17. The method of claim 14, wherein said mammal is a human.

20 18. The method of claim 14, wherein said compound inhibits translation of HO-1 mRNA in a vascular cell of said mammal.

19. The method of claim 18, wherein said compound consists of a single stranded nucleic acid complementary to at least a portion of said HO-1 mRNA.

- 34 -

20. A method of inhibiting vascular restenosis in a mammal comprising identifying a mammal having vascular restenosis or at risk of developing vascular restenosis and administering to said mammal a compound which
5 inhibits expression of HO-1.

21. The method of claim 14, wherein said compound is administered to said mammal at least one month after a vascular injury.


22. The method of claim 14, wherein said compound
10 is administered to said mammal at least two months after a vascular injury.

23. The method of claim 14, wherein said compound is administered to said mammal at least three months after a vascular injury.

15 24. A method of inhibiting vascular smooth muscle cell proliferation in a mammal comprising administering to an injured vascular tissue of said mammal a HO, wherein said HO is administered within 24 hours after a vascular injury.

20 25. The method of claim 24, wherein said HO is administered to said mammal for up to one week after a vascular injury.

FIG. 1






FIG. 2

FIG. 3





FIG. 4








FIG. 6



7/10

FIG. 7





FIG. 8



9/10

	TG (Line 451)			ZT		
MOUSE	Ventricle	spleen	Liver	Ventricle	spleen	Liver
5512				5514		
5512				5514		
5512				5514		





FIG. 10



BEST AVAILABLE COPY

SEQUENCE LISTING

<110> The President and Fellows of Harvard College

<120> INHIBITING CARDIOMYOCYTE DEATH

<130> 00246/235WO2

<150> US 60/121,946

<151> 1999-02-25

<150> US 60/098,377

<151> 1998-08-28

<160> 6

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 1550

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (81)...(944)

<400> 1

tcaacgcctg cctccccctcg agcgtcctca ggcagccgc cgcccgcgga gccagcacga	60
acgagcccag caccggcccg atg gag cgt ccg caa ccc gac agc atg ccc cag	113
Met Glu Arg Pro Gln Pro Asp Ser Met Pro Gln	
1 5 10	
gat ttg tca gag gcc ctg aag gag gcc acc aag gag gtg cac acc cag	161
Asp Leu Ser Glu Ala Leu Lys Glu Ala Thr Lys Glu Val His Thr Gln	
15 20 25	
gca gag aat gct gag ttc atg agg aac ttt cag aag ggc cag gtg acc	209
Ala Glu Asn Ala Glu Phe Met Arg Asn Phe Gln Lys Gly Gln Val Thr	
30 35 40	
cga gac ggc ttc aag ctg gtg atg gcc tcc ctg tac cac atc tat gtg	257
Arg Asp Gly Phe Lys Leu Val Met Ala Ser Leu Tyr His Ile Tyr Val	
45 50 55	
gcc ctg gag gag gag att gag cgc aac aag gag agc cca gtc ttc gcc	305
Ala Leu Glu Glu Glu Ile Glu Arg Asn Lys Glu Ser Pro Val Phe Ala	
60 65 70 75	
cct gtc tac ttc cca gaa gag ctg cac cgc aag gct gcc ctg gag cag	353
Pro Val Tyr Phe Pro Glu Glu Leu His Arg Lys Ala Ala Leu Glu Gln	
80 85 90	
gac ctg gcc ttc tgg tac ggg ccc cgc tgg cag gag gtc atc ccc tac	401
Asp Leu Ala Phe Trp Tyr Gly Pro Arg Trp Gln Glu Val Ile Pro Tyr	
95 100 105	
aca cca gcc atg cag cgc tat gtg aag cgg ctc cac gag gtg ggg cgc	449
Thr Pro Ala Met Gln Arg Tyr Val Lys Arg Leu His Glu Val Gly Arg	
110 115 120	
aca gag ccc gag ctg ctg gtg gcc cac gcc tac acc cgc tac ctg ggt	497
Thr Glu Pro Glu Leu Leu Val Ala His Ala Tyr Thr Arg Tyr Leu Gly	
125 130 135	

gac ctg tct ggg ggc cag gtg ctc aaa aag att gcc cag aaa gcc ctg 545
 Asp Leu Ser Gly Gly Gln Val Leu Lys Lys Ile Ala Gln Lys Ala Leu
 140 145 150 155
 gac ctg ccc agc tct ggc gag ggc ctg gcc ttc ttc acc ttc ccc aac 593
 Asp Leu Pro Ser Ser Gly Glu Gly Leu Ala Phe Phe Thr Phe Pro Asn
 160 165 170
 att gcc agt gcc acc aag ttc aag cag ctc tac cgc tcc cgc atg aac 641
 Ile Ala Ser Ala Thr Lys Phe Lys Gln Leu Tyr Arg Ser Arg Met Asn
 175 180 185
 tcc ctg gag atg act ccc gca gtc agg cag agg gtg ata gaa gag gcc 689
 Ser Leu Glu Met Thr Pro Ala Val Arg Gln Arg Val Ile Glu Glu Ala
 190 195 200
 aag act gcg ttc ctg ctc aac atc cag ctc ttt gag gag ttg cag gag 737
 Lys Thr Ala Phe Leu Leu Asn Ile Gln Leu Phe Glu Glu Leu Gln Glu
 205 210 215
 ctg ctg acc cat gac acc aag gac cag agc ccc tca cgg gca cca ggg 785
 Leu Leu Thr His Asp Thr Lys Asp Gln Ser Pro Ser Arg Ala Pro Gly
 220 225 230 235
 ctt cgc cag cgg gcc agc aac aaa gtg caa gat tct gcc ccc gtg gag 833
 Leu Arg Gln Arg Ala Ser Asn Lys Val Gln Asp Ser Ala Pro Val Glu
 240 245 250
 act ccc aga ggg aag ccc cca ctc aac acc cgc tcc cag gct ccg ctt 881
 Thr Pro Arg Gly Lys Pro Pro Leu Asn Thr Arg Ser Gln Ala Pro Leu
 255 260 265
 ctc cga tgg gtc ctt aca ctc agc ttt ctg gtg gcg aca gtt gct gta 929
 Leu Arg Trp Val Leu Thr Leu Ser Phe Leu Val Ala Thr Val Ala Val
 270 275 280
 ggg ctt tat gcc atg tgaatgcagg catgctggct cccagggcca tgaactttgt 984
 Gly Leu Tyr Ala Met
 285

ccggtggaag gccttctttc tagagagggg attctcttgg ctggcttcct taccgtgggc 1044
 actgaaggct ttcagggcct ccagccctct cactgtgtcc ctctctctgg aaaggaggaa 1104
 ggagcctatg gcatcttccc caacgaaaag cacatccagg caatggccta aacttcagag 1164
 ggggcgaagg ggtcagccct gcccttcagc atcctcagtt cctgcagcag agcctggaag 1224
 acaccctaag atggcagctg tctcaaacct ccaaaagccc tgagtttcaa gtatccttgt 1284
 tgacacggcc gtgaccactt tccccgtggg ccatggcaat ttttacacaa acctgaaaag 1344
 atgttggtgc ttgtgttttt gtcttatttt tggttgagcc actctgttcc tggctcagcc 1404
 tcaaatgcag tattttttgt gtgttctgtt gtttttatag caggggtggg gtgggttttg 1464
 agccatgcgt ggggtggggag ggaggtgttt aacggcactg tggccttggt ctaacttttg 1524
 tgtgaaataa taaacaacat tgtctg 1550

<210> 2

<211> 288

<212> PRT

<213> Homo sapiens

<400> 2

Met Glu Arg Pro Gln Pro Asp Ser Met Pro Gln Asp Leu Ser Glu Ala
 1 5 10 15
 Leu Lys Glu Ala Thr Lys Glu Val His Thr Gln Ala Glu Asn Ala Glu
 20 25 30
 Phe Met Arg Asn Phe Gln Lys Gly Gln Val Thr Arg Asp Gly Phe Lys
 35 40 45
 Leu Val Met Ala Ser Leu Tyr His Ile Tyr Val Ala Leu Glu Glu Glu
 50 55 60
 Ile Glu Arg Asn Lys Glu Ser Pro Val Phe Ala Pro Val Tyr Phe Pro
 65 70 75 80

aag gag gcg ctg acc aag gac atg gag tat ttc ttt ggt gaa aac tgg Lys Glu Ala Leu Thr Lys Asp Met Glu Tyr Phe Phe Gly Glu Asn Trp 110 115 120	447
gag gag cag gtg cag tgc ccc aag gct gcc cag aag tac gtg gag cgg Glu Glu Gln Val Gln Cys Pro Lys Ala Ala Gln Lys Tyr Val Glu Arg 125 130 135	495
atc cac tac ata ggg cag aac gag ccg gag cta ctg gtg gcc cat gca Ile His Tyr Ile Gly Gln Asn Glu Pro Glu Leu Leu Val Ala His Ala 140 145 150	543
tac acc cgc tac atg ggg gat ctc tcg ggg ggc cag gtg ctg aag aag Tyr Thr Arg Tyr Met Gly Asp Leu Ser Gly Gly Gln Val Leu Lys Lys 155 160 165	591
gtg gcc cag cga gca ctg aaa ctc ccc agc aca ggg gaa ggg acc cag Val Ala Gln Arg Ala Leu Lys Leu Pro Ser Thr Gly Glu Gly Thr Gln 170 175 180 185	639
ttc tac ctg ttt gag aat gtg gac aat gcc cag cag ttc aag cag ctc Phe Tyr Leu Phe Glu Asn Val Asp Asn Ala Gln Gln Phe Lys Gln Leu 190 195 200	687
tac cgg gcc agg atg aac gcc ctg gac ctg aac atg aag acc aaa gag Tyr Arg Ala Arg Met Asn Ala Leu Asp Leu Asn Met Lys Thr Lys Glu 205 210 215	735
agg atc gtg gag gcc aac aag gct ttt gag tat aac atg cag ata ttc Arg Ile Val Glu Ala Asn Lys Ala Phe Glu Tyr Asn Met Gln Ile Phe 220 225 230	783
aat gaa ctg gac cag gcc ggc tcc aca ctg gcc aga gag acc ttg gag Asn Glu Leu Asp Gln Ala Gly Ser Thr Leu Ala Arg Glu Thr Leu Glu 235 240 245	831
gat ggg ttc cct gta cac gat ggg aaa gga gac atg cgt aaa tgc cct Asp Gly Phe Pro Val His Asp Gly Lys Gly Asp Met Arg Lys Cys Pro 250 255 260 265	879
ttc tac gct gct gaa caa gac aaa ggg ctg gag ggc agc ctg tcc ctt Phe Tyr Ala Ala Glu Gln Asp Lys Gly Leu Glu Gly Ser Leu Ser Leu 270 275 280	927
ccg aca agc tat gct gtg ctg agg aag ccc agc ctc cag ttc atc ctg Pro Thr Ser Tyr Ala Val Leu Arg Lys Pro Ser Leu Gln Phe Ile Leu 285 290 295	975
gcc gct ggt gtg gcc cta gct gct gga ctc ttg gcc tgg tac tac atg Ala Ala Gly Val Ala Leu Ala Ala Gly Leu Leu Ala Trp Tyr Tyr Met 300 305 310	1023
tgaagcacc atcatgccac accggtaccc tctctccgac tgaccactgg cctaccacctt tctccagccc tgactaaact accacctcag gtgacttttt aaaaaatgct gggtttaaga aaggcaacca ataaaagaga tgctagagcc tcgtctgaca gcacacctctc tatgggccat attccgcact gggcacaggc cgtcacccctg ggagcagtcg gcacagtgca gcaagcctgg cccccgaccc agctctactc caggcttcca cacttctggg ccctaggctg cttccggtag tccctgtttt tgcagtacat ggggtgactat ctccccctgtt ggaggtgagt ggcctgtaag tccaagctgt gcgagggggc cttgctggat gctgctgtac aacttctggg cctctcttgg accctgggag tgagggtggg tgtgggtgga agcctcagag gccttgggag ctcacccctc tcacccagaa tccctctaac ccttgggtgc ggtttgctca gccccagctt atctcctcct ccgcctgtgt aaatgctcca gcaactcaata aagtgggctt tgcaagctaa aaaaaaaaaa aaaa	1083 1143 1203 1263 1323 1383 1443 1503 1563 1623 1627

<210> 4
<211> 313
<212> PRT

<213> Homo sapiens

<400> 4

```

Met Ser Ala Glu Val Glu Thr Ser Glu Gly Val Asp Glu Ser Glu Lys
 1          5          10          15
Lys Asn Ser Gly Ala Leu Glu Lys Glu Asn Gln Met Arg Met Ala Asp
          20          25          30
Leu Ser Glu Leu Leu Lys Glu Gly Thr Lys Glu Ala His Asp Arg Ala
          35          40          45
Glu Asn Thr Gln Phe Val Lys Asp Phe Leu Lys Gly Asn Ile Lys Lys
          50          55          60
Glu Leu Phe Lys Leu Ala Thr Thr Ala Leu Tyr Phe Thr Tyr Ser Ala
          65          70          75          80
Leu Glu Glu Glu Met Glu Arg Asn Lys Asp His Pro Ala Phe Ala Pro
          85          90          95
Leu Tyr Phe Pro Met Glu Leu His Arg Lys Glu Ala Leu Thr Lys Asp
          100          105          110
Met Glu Tyr Phe Phe Gly Glu Asn Trp Glu Glu Gln Val Gln Cys Pro
          115          120          125
Lys Ala Ala Gln Lys Tyr Val Glu Arg Ile His Tyr Ile Gly Gln Asn
          130          135          140
Glu Pro Glu Leu Leu Val Ala His Ala Tyr Thr Arg Tyr Met Gly Asp
          145          150          155          160
Leu Ser Gly Gly Gln Val Leu Lys Lys Val Ala Gln Arg Ala Leu Lys
          165          170          175
Leu Pro Ser Thr Gly Glu Gly Thr Gln Phe Tyr Leu Phe Glu Asn Val
          180          185          190
Asp Asn Ala Gln Gln Phe Lys Gln Leu Tyr Arg Ala Arg Met Asn Ala
          195          200          205
Leu Asp Leu Asn Met Lys Thr Lys Glu Arg Ile Val Glu Ala Asn Lys
          210          215          220
Ala Phe Glu Tyr Asn Met Gln Ile Phe Asn Glu Leu Asp Gln Ala Gly
          225          230          235          240
Ser Thr Leu Ala Arg Glu Thr Leu Glu Asp Gly Phe Pro Val His Asp
          245          250          255
Gly Lys Gly Asp Met Arg Lys Cys Pro Phe Tyr Ala Ala Glu Gln Asp
          260          265          270
Lys Gly Leu Glu Gly Ser Leu Ser Leu Pro Thr Ser Tyr Ala Val Leu
          275          280          285
Arg Lys Pro Ser Leu Gln Phe Ile Leu Ala Ala Gly Val Ala Leu Ala
          290          295          300
Ala Gly Leu Leu Ala Trp Tyr Tyr Met
          305          310

```

<210> 5

<211> 2225

<212> DNA

<213> Rattus rattus

<220>

<221> CDS

<222> (1062)...(1931)

<400> 5

```

tttcagggat ttttgcgatt cctctctgta gacttctact tgttctctaa gggagttctt 60
catgtctttc ttgaagtcac ccagcatcat gatcaaatat gattttgaaa ctagatcttg 120
cttttctggt gtgtttggat attccatgtt tgttttggtg ggagaattgg gctccgatga 180
tggcattgtag tcttggtttc tgttgcttgg tttcctgccc ttgcctctcg ccatcagatt 240
atctctagtg ttactttggt ctgctatttc tgacagtggc tagactgtcc tataagcctg 300
tgtgtcagga gtgctgtaga ccttttttcc tctctttcag tcagttatgg gacagagtgt 360
tctgcttttg ggcgtgtagt ttttcctctc tacaggtctt cagctgttcc tgtgggcctg 420
tgtcttgagt tcaccaggca gctttcttgc agcagaaaaa ttgggtcatc ctgtgatcct 480
gaggctcaag ttoctctcgt gggtgctgtc caggggctct ctgcagcggg cacaaccagg 540
aagacctgtg cggccccctc cggagcttca gtgcaccagg gttccagatg gcctttggcg 600
ttttcctctg gcgtccgaga tgtatgtaca gagagcagtc tcttctgggt tcccaggctt 660
gtctgcctct ctgaagggtc agctctccct cccacgggat ttgggtgcag agaactgttt 720
atccggtctg tttctttcag gttccggtgg tgtctcagga aggtgtcgtt cctgcgcctt 780

```

cccccatggg	accagaggcc	ttatacagtt	tcctcttggg	ccagggatgt	gggcaggggt	840
gagcagtgtt	ggtggtctct	tccgtctgca	gcctcaggag	tgccacctga	ccaggcggtt	900
gggtctctct	ctgagaattt	catttttaaa	tcattcatta	aaatgtcatg	acttgatgtc	960
ctgctgtccg	tctcacgccc	tcagctgtaa	cagtgccgag	ggagtcaactg	aagaagagac	1020
tgaatgacca	gagtatgggc	agcacagaca	actcaacaaa	a atg tct tca gag gtg		1076
				Met Ser Ser Glu Val		
				1	5	
gag act gcg gag gcc gta gat gag tca gag aag aac tct atg gca tca						1124
Glu Thr Ala Glu Ala Val Asp Glu Ser Glu Lys Asn Ser Met Ala Ser						
	10		15		20	
gag aag gaa aac cat tcc aaa ata gca gac ttt tct gat ctt ctg aag						1172
Glu Lys Glu Asn His Ser Lys Ile Ala Asp Phe Ser Asp Leu Leu Lys						
	25		30		35	
gaa ggg aca aag gaa gca gat gac cgg gca gaa aat acc cag ttt gtc						1220
Glu Gly Thr Lys Glu Ala Asp Asp Arg Ala Glu Asn Thr Gln Phe Val						
	40		45		50	
aaa gac ttc ttg aaa gga aac att aag aag gag cta ttt aag ctg gcc						1268
Lys Asp Phe Leu Lys Gly Asn Ile Lys Lys Glu Leu Phe Lys Leu Ala						
	55		60		65	
acc act gca ctt tca tac tca gcc cct gag gag gaa atg gat tca ctg						1316
Thr Thr Ala Leu Ser Tyr Ser Ala Pro Glu Glu Glu Met Asp Ser Leu						
	70		75		80	85
acc aag gac atg gag tac ttc ttt ggt gaa aac tgg gag gaa aaa gtg						1364
Thr Lys Asp Met Glu Tyr Phe Phe Gly Glu Asn Trp Glu Glu Lys Val						
	90		95		100	
aag tgc tct gaa gct gcc cag acg tat gtg gat cag att cac tat gta						1412
Lys Cys Ser Glu Ala Ala Gln Thr Tyr Val Asp Gln Ile His Tyr Val						
	105		110		115	
ggg caa aat gag cca gag cat ctg gtg gcc cat act tac tct act tac						1460
Gly Gln Asn Glu Pro Glu His Leu Val Ala His Thr Tyr Ser Thr Tyr						
	120		125		130	
atg ggg gga aac ctt tca ggg gac cag gta ctg aag aag gag acc cag						1508
Met Gly Gly Asn Leu Ser Gly Asp Gln Val Leu Lys Lys Glu Thr Gln						
	135		140		145	
ccg gtc ccc ttc act agg gaa ggg act cag ttc tac ctg ttt gag cat						1556
Pro Val Pro Phe Thr Arg Glu Gly Thr Gln Phe Tyr Leu Phe Glu His						
	150		155		160	165
gta gac aat gct aag caa ttc aag cta ttc tac tgc gct aga ttg aat						1604
Val Asp Asn Ala Lys Gln Phe Lys Leu Phe Tyr Cys Ala Arg Leu Asn						
	170		175		180	
gcc ttg gac ctg aat ttg aag acc aaa gag agg att gtg gag gaa gcc						1652
Ala Leu Asp Leu Asn Leu Lys Thr Lys Glu Arg Ile Val Glu Glu Ala						
	185		190		195	
acc aaa gcc ttt gaa tat aat atg cag ata ttc agt gaa ctg gac cag						1700
Thr Lys Ala Phe Glu Tyr Asn Met Gln Ile Phe Ser Glu Leu Asp Gln						
	200		205		210	
gca ggc tcc ata cca gta aga gaa acc cta aag aat ggg ctc tca ata						1748
Ala Gly Ser Ile Pro Val Arg Glu Thr Leu Lys Asn Gly Leu Ser Ile						
	215		220		225	
ctt gat ggg aag gga ggt gta tgc aaa tgt ccc ttt aat gct gct cag						1796
Leu Asp Gly Lys Gly Gly Val Cys Lys Cys Pro Phe Asn Ala Ala Gln						
	230		235		240	245

cca gac aaa ggt acc ctg gga ggc agc aac tgc cct ttc cag atg tcc 1844
Pro Asp Lys Gly Thr Leu Gly Gly Ser Asn Cys Pro Phe Gln Met Ser 260
250 255

atg gcc ttg ctg agg aag cct aac ttg cag ctc att cta gtt gcc agt 1892
Met Ala Leu Leu Arg Lys Pro Asn Leu Gln Leu Ile Leu Val Ala Ser 275
265 270

atg gcc ttg gta gct gga ctt tta gcc tgg tac tac atg tgaagggcct 1941
Met Ala Leu Val Ala Gly Leu Leu Ala Trp Tyr Tyr Met 290
280 285

gtcaagttgt ttgcatccta tctcaacatc ctaccacttg ttccttcccc acctccacct 2001
ctgcctagaa ctaccacctc aggtgacatt tttaatgttg ggtttgagaa aatgagcaac 2061
caataaaaga cagaccctag aaaaaagtca tgacttaagt ggcacgggga cacctaaagt 2121
cacactttgt gcttcagaca tactttcttt ctctatttca acactgaatt cggaagtaa 2181
cctactacta ttaataataa atgctacaca atgcataata aaaa 2225

<210> 6
<211> 290
<212> PRT
<213> Rattus rattus

<400> 6
Met Ser Ser Glu Val Glu Thr Ala Glu Ala Val Asp Glu Ser Glu Lys 15
1 5 10
Asn Ser Met Ala Ser Glu Lys Glu Asn His Ser Lys Ile Ala Asp Phe 30
20 25
Ser Asp Leu Leu Lys Glu Gly Thr Lys Glu Ala Asp Asp Arg Ala Glu 45
35 40
Asn Thr Gln Phe Val Lys Asp Phe Leu Lys Gly Asn Ile Lys Lys Glu 60
50 55
Leu Phe Lys Leu Ala Thr Thr Ala Leu Ser Tyr Ser Ala Pro Glu Glu 80
65 70
Glu Met Asp Ser Leu Thr Lys Asp Met Glu Tyr Phe Phe Gly Glu Asn 95
85 90
Trp Glu Glu Lys Val Lys Cys Ser Glu Ala Ala Gln Thr Tyr Val Asp 110
100 105
Gln Ile His Tyr Val Gly Gln Asn Glu Pro Glu His Leu Val Ala His 125
115 120
Thr Tyr Ser Thr Tyr Met Gly Gly Asn Leu Ser Gly Asp Gln Val Leu 140
130 135
Lys Lys Glu Thr Gln Pro Val Pro Phe Thr Arg Glu Gly Thr Gln Phe 160
145 150
Tyr Leu Phe Glu His Val Asp Asn Ala Lys Gln Phe Lys Leu Phe Tyr 175
165 170
Cys Ala Arg Leu Asn Ala Leu Asp Leu Asn Leu Lys Thr Lys Glu Arg 190
180 185
Ile Val Glu Glu Ala Thr Lys Ala Phe Glu Tyr Asn Met Gln Ile Phe 205
195 200
Ser Glu Leu Asp Gln Ala Gly Ser Ile Pro Val Arg Glu Thr Leu Lys 220
210 215
Asn Gly Leu Ser Ile Leu Asp Gly Lys Gly Gly Val Cys Lys Cys Pro 240
225 230
Phe Asn Ala Ala Gln Pro Asp Lys Gly Thr Leu Gly Gly Ser Asn Cys 255
245 250
Pro Phe Gln Met Ser Met Ala Leu Leu Arg Lys Pro Asn Leu Gln Leu 270
260 265
Ile Leu Val Ala Ser Met Ala Leu Val Ala Gly Leu Leu Ala Trp Tyr 285
275 280
Tyr Met 290

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 38/44, 48/00, A61P 9/10 // A61K 38/18, 31/555		A3	(11) International Publication Number: WO 00/12118
			(43) International Publication Date: 9 March 2000 (09.03.00)
(21) International Application Number: PCT/US99/19823 (22) International Filing Date: 27 August 1999 (27.08.99) (30) Priority Data: 60/098,377 28 August 1998 (28.08.98) US 60/121,946 25 February 1999 (25.02.99) US (71) Applicant: PRESIDENT AND FELLOWS OF HARVARD COLLEGE [US/US]; 17 Quincy Street, Cambridge, MA 02138 (US). (71)(72) Applicants and Inventors: LEE, Mu-En [CN/US]; 102 Nardell Road, Newton, MA 02159 (US). PERRELLA, Mark, A. [US/US]; 33 Pond Avenue, #420, Brookline, MA 02146 (US). YET, Shaw-Fang [CN/US]; 9 Donald Circle, Andover, MA 01810 (US). (74) Agent: BEATTIE, Ingrid, A.; Fish & Richardson P.C., 225 Franklin Street, Boston, MA 02110-2804 (US).			(81) Designated States: AU, CA, IL, JP, NO, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> (88) Date of publication of the international search report: 29 June 2000 (29.06.00)
(54) Title: INHIBITING CARDIOMYOCYTE DEATH			
(57) Abstract The invention features methods of inhibiting cardiomyocyte death in a mammal by administering to the myocardium of the mammal a heme oxygenase (HO). Methods of preserving an organ for transplantation and methods of inhibiting vascular stenosis are also within the invention.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

Application No

PCT/US 99/19823

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K38/44 A61K48/00 A61P9/10 //A61K38/18,A61K31/555

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ABRAHAM, N. G. (1) ET AL: "Transfection of the human heme oxygenase gene into rabbit coronary microvessel endothelial cells: Protective effect against heme and hemoglobin toxicity." PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, (1995) VOL. 92, NO. 15, PP. 6798-6802. , XP002100374 the whole document ---	1-12
X	WO 98 08566 A (WISCONSIN MED COLLEGE INC ;UNIV DUKE (US)) 5 March 1998 (1998-03-05) page 4, line 23 -page 6, line 12 page 7, line 24 -page 8, line 2 page 9, line 19 -page 10, line 18 page 55, line 2 - line 30 page 57, line 22 - line 28 --- -/--	14-25

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

23 February 2000

Date of mailing of the international search report

09/03/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Stein, A

INTERNATIONAL SEARCH REPORT

Application No

PCT/US 99/19823

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WO 97 36615 A (HARVARD COLLEGE) 9 October 1997 (1997-10-09) page 2, line 11 - line 36 page 7, line 16 -page 8, line 13 page 16, line 18 -page 17, line 36 page 21, line 14 -page 22, line 8 claims 1-6</p>	14-23
A	<p>MAULIK N ET AL: "Nitric oxide/carbon monoxide. A molecular switch for myocardial preservation during ischemia." CIRCULATION, (1996 NOV 1) 94 (9 SUPPL) II398-406. , XP000876907 the whole document</p>	1-13
A	<p>ABRAHAM, NADER G. (1): "Manipulation of heme oxygenase expression by gene transfer and metals: Implications in cell injury and repair." JOURNAL OF NEUROCHEMISTRY, (1998) VOL. 70, NO. SUPPL. 1, PP. S45. MEETING INFO.: 29TH ANNUAL MEETING OF THE AMERICAN SOCIETY FOR NEUROCHEMISTRY DENVER, COLORADO, USA MARCH 7-11, 1998 AMERICAN SOCIETY FOR NEUROCHEMISTRY. , XP000876935 the whole document</p>	1-13
A	<p>LONG, XILIN ET AL: "Hypoxia-induced expression of heme oxygenase gene expression in cultured neonatal rat cardiac myocytes." CIRCULATION, (1995) VOL. 92, NO. 8 SUPPL., PP. I653-I654, XP000876926 the whole document</p>	1-13
A	<p>HOSHIDA, SHIRO ET AL: "Heme oxygenase -1 as a culture shock protein in rat neonatal cardiomyocytes." JOURNAL OF MOLECULAR AND CELLULAR CARDIOLOGY, (1994) VOL. 26, NO. 11, PP. CCXII, XP000876927 the whole document</p>	1-13
A	<p>BORGER DR: "Induction of HSP 32 gene in hypoxic cardiomyocytes is attenuated by treatment with N-acetyl-L-cysteine" AMERICAN JOURNAL OF PHYSIOLOGY, vol. 274, no. 3 Pt 2, March 1998 (1998-03), pages H965-73, XP002131421 the whole document</p>	1-14

-/--

INTERNATIONAL SEARCH REPORT

Application No
PCT/US 99/19823

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	MORITA T ET AL: "Carbon monoxide controls the proliferation of hypoxic vascular smooth muscle cells" JOURNAL OF BIOLOGICAL CHEMISTRY, (26 DEC 1997) VOL. 272, NO. 52, PP. 32804-32809, XP002131422 the whole document	24,25
P,X	SOARES M P ET AL: "Expression of heme oxygenase -1 can determine cardiac xenograft survival." NATURE MEDICINE, (1998 SEP) 4 (9) 1073-7., XP002131423 the whole document	13

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/ 19823

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 1-8, 14-25 are directed to a method of treatment of the human/animal the search has been carried out and based on the alleged effects of the compound /composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Application No

PCT/US 99/19823

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9808566 A	05-03-1998	AU 4054297 A EP 0963219 A	19-03-1998 15-12-1999
WO 9736615 A	09-10-1997	US 5888982 A	30-03-1999



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 38/44, 48/00, A61P 9/10 // A61K 38/18, 31/555		Á3	(11) International Publication Number: WO 00/12118
			(43) International Publication Date: 9 March 2000 (09.03.00)
(21) International Application Number: PCT/US99/19823		(81) Designated States: AU, CA, IL, JP, NO, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 27 August 1999 (27.08.99)			
(30) Priority Data: 60/098,377 28 August 1998 (28.08.98) US 60/121,946 25 February 1999 (25.02.99) US		Published <i>With international search report.</i>	
(71) Applicant: PRESIDENT AND FELLOWS OF HARVARD COLLEGE [US/US]; 17 Quincy Street, Cambridge, MA 02138 (US).		(88) Date of publication of the international search report: 29 June 2000 (29.06.00)	
(72) Inventors: LEE, Mu-En; 102 Nardell Road, Newton, MA 02159 (US). PERRELLA, Mark, A.; 33 Pond Avenue #420, Brookline, MA 02146 (US). YET, Shaw-Fang; 9 Donald Circle, Andover, MA 01810 (US).			
(74) Agent: BEATTIE, Ingrid, A.; Fish & Richardson P.C., 225 Franklin Street, Boston, MA 02110-2804 (US).			
(54) Title: INHIBITING CARDIOMYOCYTE DEATH			
(57) Abstract			
<p>The invention features methods of inhibiting cardiomyocyte death in a mammal by administering to the myocardium of the mammal a heme oxygenase (HO). Methods of preserving an organ for transplantation and methods of inhibiting vascular stenosis are also within the invention.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**CORRECTED
VERSION***

**CORRECTED
VERSION****

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 38/44, 48/00, A61P 9/10 // A61K 38/18, 31/555		Ä3	(11) International Publication Number: WO 00/12118
			(43) International Publication Date: 9 March 2000 (09.03.00)
(21) International Application Number: PCT/US99/19823		(81) Designated States: AU, CA, IL, JP, NO, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 27 August 1999 (27.08.99)		Published <i>With international search report.</i>	
(30) Priority Data: 60/098,377 28 August 1998 (28.08.98) US 60/121,946 25 February 1999 (25.02.99) US		(88) Date of publication of the international search report: 29 June 2000 (29.06.00)	
(71) Applicant: PRESIDENT AND FELLOWS OF HARVARD COLLEGE [US/US]; 17 Quincy Street, Cambridge, MA 02138 (US).			
(72) Inventors: LEE, Mu-En; 102 Nardell Road, Newton, MA 02159 (US). PERRELLA, Mark, A.; 33 Pond Avenue #420, Brookline, MA 02146 (US). YET, Shaw-Fang; 9 Donald Circle, Andover, MA 01810 (US).			
(74) Agent: BEATTIE, Ingrid, A.; Fish & Richardson P.C., 225 Franklin Street, Boston, MA 02110-2804 (US).			
(54) Title: INHIBITING CARDIOMYOCYTE DEATH			
(57) Abstract The invention features methods of inhibiting cardiomyocyte death in a mammal by administering to the myocardium of the mammal a heme oxygenase (HO). Methods of preserving an organ for transplantation and methods of inhibiting vascular stenosis are also within the invention.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

- 1 -

INHIBITING CARDIOMYOCYTE DEATHRelated Application Information

5 This application claims priority from provisional application no. 60/121,946, filed on February 25, 1999, and provisional application no. 60/098,377, filed on August 28, 1998.

Statement as to Federally Sponsored Research

10 This invention was made with U.S. Government support under National Institutes of Health grants RO1 GM53249, KO8 HL03274, and KO8 HL03194. The government has certain rights in the invention.

Background of the Invention

15 The invention relates to treatment of cardiovascular disease.

Myocardial infarction is one of the most common diagnoses of hospitalized patients in western countries. In the United States, over 1.5 million myocardial infarctions occur annually, and mortality from acute myocardial infarction is approximately 25 per cent. Thrombolytic therapy and reperfusion of ischemic myocardium, e.g., using percutaneous transluminal coronary angioplasty (PTCA), have decreased mortality rates among patients who have suffered an acute myocardial infarction. Nevertheless, myocardial damage and heart failure resulting from a failure to achieve early revascularization or from reperfusion injury remain a significant clinical problem.

Summary of the Invention

30 The invention features methods of minimizing myocardial damage by salvaging hypoxic myocardial tissue before it becomes irreversibly injured. For example, a method of inhibiting cardiomyocyte death in a mammal, e.g., a human, who has suffered a myocardial infarction or who has myocarditis is carried out by locally

- 2 -

administering to the myocardium of the mammal a heme oxygenase (HO) polypeptide. Preferably, the HO polypeptide has the amino acid sequence of a naturally-occurring heme oxygenase-1 (HO-1), heme oxygenase-2 (HO-2), or heme oxygenase-3 (HO-3), or a biologically active fragment thereof.

Compositions, such as hemin, hemoglobin, or heavy metals, e.g., tin or nickel, that increase production of endogenous HO, are also administered to inhibit cardiomyocyte death or damage. For example, overexpression of HO-1 is induced in vascular cells by exposure to heme, heavy metals, endotoxin and hyperoxia, hyperthermia, shear stress and strain, UV light, or reactive oxygen species. By "overexpression" is meant a level of protein production that at least 20% greater than that present in the tissue under normal physiologic conditions. Preferably, the level of HO-1 expression in vascular tissue in the presence of an inducing agent is at least 20% greater than that in the absence of the inducing agent; more preferably, the level of expression is at least 50% greater, more preferably, the level of expression is at least 100% greater, and most preferably, the level of expression is at least 200% greater than that in the absence of an inducing agent. Inhibition of cardiomyocyte death is also achieved by locally administering to the myocardium of a mammal a DNA encoding a HO. HO expression by target cell, e.g., VSMC, is increased by administering to the cells exogenous DNA encoding HO, e.g., a plasmid containing DNA encoding human HO-1 or HO-2 under the control of a strong constitutive promoter. Oxidative stress leads to cell death by apoptosis and/or necrosis. By neutralizing reactive oxygen species, HO reduces cardiomyocyte damage and death due to oxidative stress.

- 3 -

The invention also includes a method of inhibiting cardiomyocyte death *in vitro* by contacting cardiomyocytes with an HO or DNA encoding an HO. For example, a method of preserving isolated myocardial tissue, e.g., a donor heart to be used for transplantation, is carried out by bathing or perfusing the tissue with a solution containing an HO or a DNA encoding an HO. The method allows prolonged storage of organs after removal from the donor and prior to transplantation into a recipient by reducing irreversible ischemic tissue damage. By "isolated myocardial tissue" is meant tissue that has been removed from a living or recently deceased mammal. Preferably, a donor heart is preserved in an HO solution for 0.5-6 hours prior to transplantation. More preferably, the organ is preserved for greater than 6 hours, e.g., 8, 10, 12, and up to 24 hours.

A method of inhibiting vascular stenosis or restenosis in a mammal, e.g., a human, is also within the invention. The method is carried out by locally administering to the site of a vascular injury or a site which is at risk of developing a stenotic lesion a compound which inhibits expression of HO-1, and as a result, VSMC proliferation. To inhibit vascular stenosis or restenosis (which occurs at a relatively late stage after the occurrence of a vascular injury), the compound is administered at least one month after an injury such as surgery or angioplasty. For example, such treatment is administered 3 weeks to several months (e.g., 2 months or 3 months) post-injury. Preferably, the compound inhibits transcription of the gene encoding HO-1 or inhibits translation of HO-1 mRNA into an HO-1 polypeptide in a vascular cell, e.g., a vascular smooth muscle cell (VSMC), of the mammal. The vascular cell is preferably an aortic smooth muscle cell, e.g., an aortic smooth muscle cell located in the region of an artery

- 4 -

affected by vascular stenosis or restenosis such as the site of balloon angioplasty or coronary bypass surgery. For example, transforming growth factor- β 1 (TGF- β 1) is administered to inhibit production of HO-1 mRNA and HO gene product.

For antisense therapy, the compound is a antisense nucleic acid molecule containing at least 10 nucleotides, the sequence of which is complementary to an mRNA encoding all or part of a wild type HO polypeptide.

10 Preferably, the compound, e.g., an antisense oligonucleotide or antisense RNA produced from an antisense template, inhibits HO expression. The antisense nucleic acid inhibits HO expression by inhibiting translation of HO mRNA. For example,

15 antisense therapy is carried out by administering a single stranded nucleic acid complementary at least a portion of HO mRNA to interfere with the translation of mRNA into protein, thus reducing the amount of functional HO produced in the cell. The method includes the step of

20 identifying a mammal having undesired vascular stenosis or restenosis or at risk of developing such a condition. For example, the mammal to be treated is one who needs or has recently undergone PTCA, coronary artery bypass surgery, other vascular injury, that stimulates vascular

25 smooth muscle cell proliferation that results in undesired vascular stenosis or restenosis.

For treatment of a vascular injury soon after the injury or stress has occurred, an HO polypeptide, e.g., HO-1, or a nucleic acid encoding an HO polypeptide, is

30 administered to a mammal within minutes until approximately one week post-injury. Augmentation of the level of HO in injured vascular tissue shortly after the injury has occurred inhibits an initial increase in local VSMC proliferation post-injury. For example, such early

- 5 -

stage intervention is carried out within 24 hours post-injury.

Other features and advantages of the invention will be apparent from the following detailed description,
5 and from the claims.

Detailed Description

The drawings will first be briefly described.

Drawings

Fig. 1 is a diagram of the targeted gene
10 disruption strategy used in making an HO-1-deficient mouse.

Fig. 2 is a bar graph showing that hypoxia increases hematocrit in HO-1 +/+ and -/- mice.

Fig. 3 is a bar graph showing that hypoxia
15 markedly increases ventricular weight in HO-1 -/- mice.

Fig. 4 is a diagram of the mouse model of vein graft stenosis.

Fig. 5A is a line graph showing luminal occlusion of an artery into which a vein patch has been grafted.

20 Fig. 5B is a diagram of a vein graft.

Fig. 6 is a diagram of plasmid containing a myosin heavy chain promoter which directs cardiospecific expression of a polypeptide-encoding DNA to which it is operably linked.

25 Fig. 7 is a bar graph showing that chronic hypoxia increases right ventricular systolic pressure. Wild type (+/+) and HO-1-deficient (-/-) mice were exposed to normoxia or chronic hypoxia (10% oxygen). Right ventricular pressure was measured under normoxic
30 conditions (open bars) or after five weeks of hypoxia (filled bars). Error bars indicate standard deviation. *P<0.05 vs. animals exposed to normoxia within the same group (n = 5 in each group).

Fig. 8 is a bar graph showing that HO-1 -/-
35 arterial smooth muscle cells are more sensitive to

- 6 -

oxidative stress compared to wild type smooth muscle cells.

Fig. 9 is an autoradiograph of a Northern blot assay showing expression of a human HO-1 (hHO-1)

5 transgene in a transgenic mouse.

Fig. 10 is an autoradiograph of a Western blot showing the presence of a hHO-1 gene product in tissues of HO-1 transgenic mice.

HO-1-deficient mice

10 HO-1-deficient (HO-1-/-) mice were produced using a standard targeted gene deletion strategy to delete exon 3 (Fig. 1). The murine HO-1 gene contains 5 exons and 4 introns, spanning approximately 7 kilobases (kb). The targeting construct was made by deleting the largest exon
15 (exon 3) which contains 492 nucleotides out of the 867 nucleotides of the entire open reading frame. This deletion renders the HO-1 enzyme non-functional. An *XhoI/BamHI* fragment of the neo cassette from pMC1neo PolyA plasmid was subcloned into pBluescript II SK
20 (Stratagene, La Jolla, CA) to generate pBS-neo. To generate pBS-neo-HO-1, the 3 kb *XhoI* fragment of the HO-1 gene spanning from exon 1 to the end of intron 2 was subcloned into the *XhoI* site of pBS-neo in the same orientation as the neo cassette. The 4 kb HO-1 *BamHI*-
25 *EcoRI* fragment containing a small portion of intron 3, exon 4, and exon 5 was subcloned into *BamHI* and *EcoRI* site of pPGK-TK to generate pPGK-TK-HO-1. The 7 kb *BamHI*-*ClaI* fragment (filled in with Klenow) from pPGK-TK-HO-1 was then subcloned into *BamHI* and *XbaI* sites (filled
30 in with Klenow) sites of pBS-neo-HO-1 to generate the HO-1 targeting construct. The linearized targeting construct was transfected into murine D3 embryonic stem (ES) cells, and a clone with the correct homologous recombination (yielding the appropriately disrupted HO-1
35 gene) injected into blastocysts and used to generate HO-1

- 7 -

deficient mice. The survival rate of HO-1 $-/-$ mice was 25% of the expected survival rate, and the mice were grossly normal. The mice were deficient in HO-1 mRNA and HO-1 protein but not HO-2 mRNA or protein.

5 HO-1 transgenic mice

Standard techniques were used to generate mice which express a hHO-1 transgene in heart tissue. The transgene was cloned under the control of the cardiac α -myosin heavy chain promoter for expression preferentially
10 in cardiovascular tissue. One group of transgenic mice were engineered to express hHO-1 DNA in the sense orientation, and another group expressed hHO-1 DNA in the antisense orientation. As shown in Fig. 9, hHO-1 mRNA was detected in the heart (ventricle) of the transgenic
15 mouse but not in other tissues tested. Western blot analysis confirmed the presence of a transgenic hHO-1 gene product in heart tissue (ventricles) of hHO-1 transgenic mice. hHO-1 transgenic mice are used to evaluate the effect of HO-1 expression (and
20 overexpression) in cardiovascular tissue, e.g., in response to injury or stress.

Inhibition of cardiomyocyte death

Oxidative stress caused by such conditions as ischemia and reperfusion injury induces myocardial
25 dysfunction and cardiomyocyte death. HO is an enzyme that catalyzes oxidation of heme to generate carbon monoxide (CO; which can increase cellular cGMP) and biliverdin (which is a potent antioxidant). HO-1 is an inducible isoform of HO, whereas HO-2 is constitutively
30 expressed. Expression of HO-1 is induced in the cardiovascular system by such stimuli as hypoxia, hyperoxia, cytokines such as interleukin- 1β (IL- 1β), endotoxemia, heat shock, and ischemia. HO-1 also regulates VSMC growth. Expression of inducible HO (HO-1)

- 8 -

is markedly induced in the cardiovascular system by stimuli such as increased pressure and hypoxia.

To study the effect of HO-1 on mammalian responses to hypoxia such as that manifested in clinical conditions, e.g., high altitude pulmonary edema, myocardial infarction, myocarditis, pulmonary hypertension, pulmonary embolism, pulmonary valve stenosis, congenital heart disease, and chronic obstructive pulmonary disease (e.g., emphysema), mice were subjected to chronic hypoxia. In accordance with a standard model of pulmonary hypertension, mice were kept in a 10% O₂ chamber for 7 weeks. Two groups (Group I = five HO-1 +/+ mice; Group II = five HO-1 -/- mice) were studied. Two of the HO-1 deficient mice died at week 7; none of the HO-1 +/+ mice died. As shown in Fig. 2, exposure of the mice to hypoxic conditions resulted in an increase in hematocrit in both wild type and knockout (HO-1 -/-) mice, indicating a high level of tissue hypoxia of the mice. Under normoxia conditions, the heart weight of HO-1-deficient and wild type mice was comparable, but under hypoxic conditions the ventricular weight of HO-1-deficient mice greater than that of the HO-1-deficient mice kept under normoxic conditions (Fig. 3). For example, exposure to hypoxia for 7 weeks caused a 32% increase in the ventricular weight index in wild type mice, whereas in HO-1-deficient mice, the ventricular weight index after 7 weeks of hypoxia increased 100% (compared to normoxic wild type mice. Changes in the ventricular weight reflected mainly a right ventricular effect, as the RV(LV+septum) increased in HO-1-deficient mice compared to wild type mice exposed to hypoxia for 7 weeks.

Fig. 7 shows the effect of hypoxia on right ventricular systolic pressure, an indicator of pulmonary arterial systolic pressure. Right ventricular systolic

- 9 -

pressure in wild type and HO-1 $-/-$ mice did not differ under normoxic conditions ($P = 0.80$; Fig. 7, open bars). Although five weeks of hypoxia increased right ventricular systolic pressure, it did so to a similar degree in wild type and HO-1 $-/-$ mice ($P = 0.43$; Fig. 7, filled bars).

In HO-1-deficient mice, exposure to conditions of chronic hypoxia resulted in more dramatic hypertrophy and dilation of the right ventricle of the heart compared to that observed in wild type mice. Evidence of massive cardiomyocyte death was detected and large organized thrombi attached to areas of infarct were also detected in HO-1-deficient mice but not in wild type mice.

When stained with Masson's trichrome stain (which detects collagen), an increase in collagen fibers was observed in tissue sections of the right ventricle of HO-1 $-/-$ mice in response to hypoxia compared to the amount of collagen detected in wild type sections. These data indicate evidence of scar formation and repair mechanisms in the hearts of HO-1-deficient mice under hypoxic conditions. Tissue sections were also stained with PECAM stain (which detects endothelial cells). Increased blood vessel formation was detected in the right ventricles of HO-1 $-/-$ mice in response to hypoxia compared to wild type mice. These data suggest that HO-1 inhibits angiogenesis in response to hypoxia.

The mechanism of cardiomyocyte death in HO-1 $-/-$ mice under hypoxic conditions was evaluated by histological analysis, immunocytochemistry, and TdT-mediated dUTP-biotin nickend labeling (TUNEL assay). The standard TUNEL assay detects apoptosis. Ventricles were fixed in 4% paraformaldehyde overnight at 4°C and embedded in paraffin. Tissue sections were stained with hematoxylin and eosin or Masson's trichrome. To detect oxidation-specific lipid-protein adducts, heart tissue

- 10 -

sections were immunostained with polyclonal antibody MAL-2 (anti-malondialdehyde-lysine; Rosenfeld et al., 1990, Arteriosclerosis 10:336-349) and counterstained with methyl green. TUNEL was used to detect DNA breaks in apoptotic cells *in situ*. Red staining in the nuclei of the cells indicated a positive reaction; the cells were also counterstained with methyl green. Pulmonary vascular remodeling was assessed in lungs that had been perfused with saline through the pulmonary artery and fixed with 4% paraformaldehyde instilled through the trachea. Muscularization of peripheral vessels was determined using standard methods, e.g., that described in Klinger et al., 1993, J. Appl. Physiol. 75:198-205. The nuclei of cardiomyocytes of HO-1-deficient mice subjected to hypoxic conditions stained positive, providing evidence that apoptosis contributes to the mechanism of death of cardiomyocytes under these conditions.

Additional histological analyses were undertaken to confirm that chronic hypoxia induces right ventricular infarction in HO-1-deficient mice. Cardiomyocytes were intact in ventricular sections from wild type mice exposed to 7 weeks of hypoxia, but ventricular sections from HO-1-deficient mice exposed to 7 weeks of hypoxia showed mononuclear inflammatory cell infiltration, extensive cardiomyocyte degeneration, and death with focal calcification. These observations indicate that infarcts were 1-2 weeks old. The right ventricular infarcts did not appear to result from vascular occlusion, because the coronary arteries supplying blood to the right ventricle were patent in HO-1-deficient mice.

To detect collagen accumulation indicative of fibrosis, ventricular sections were stained with Masson's trichrome. After 7 weeks of hypoxia, cells surrounding

- 11 -

blood vessels stained positive for collagen in hearts from wild type mice. In hearts from HO-1-deficient mice exposed to hypoxia for 7 weeks, early collagen deposition was present throughout the lesion. The degree of
5 fibrosis was consistent with early scar formation and a lesion of 1-2 weeks old.

Two out of the six hearts from 7-week hypoxic HO-1-deficient mice did not exhibit right ventricular infarcts; however, close examination of the hearts
10 revealed focal areas of myocardial degeneration without evidence of extensive inflammatory cell infiltration. These hearts showed early evidence of myocardial damage. Heart from wild type and HO-1-deficient mice examined after 5 weeks of hypoxia displayed no cardiomyocyte
15 degeneration or death and no extensive mononuclear inflammatory cell infiltration. Cells surrounding blood vessels stained positive for collagen in hearts from wild type and HO-1-deficient mice housed for 5 weeks under normoxic and hypoxic conditions. In contrast with the
20 right ventricular free walls from HO-1-deficient mice exposed to 7 weeks of hypoxia (which showed deposition of collagen), no collagen deposition was evident in the right ventricular free walls from HO-1-deficient mice exposed to 5 weeks of hypoxia. These data confirm that
25 in the HO-1-deficient mice exposed to 7 weeks of hypoxia, myocardial infarcts were less than 2 weeks old.

To assess oxidative damage in hearts with infarcts taken from HO-1-deficient mice, ventricular tissue sections were immunostained with MAL-2 which detects
30 oxidation-specific lipid-protein adducts. In contrast to the minimal MAL-2 staining observed in hearts from wild type mice, MAL-2 staining was intense in cells (predominantly cardiomyocytes) beneath the infarcted area of the right ventricle in HO-1-deficient mice. These
35 data indicate the presence of severe oxidative damage

- 12 -

within and around the infarct site. No TUNEL-positive cardiomyocytes were detectable in right ventricles from wild-type mice, but a significant number of TUNEL-positive cells surrounded the infarct site in right
5 ventricles form HO-1-deficient mice.

The data described herein indicate that
(1) pulmonary vascular remodeling in response to hypoxia is similar in HO-1 +/+ and -/- mice, (2) hypoxia induces more severe right ventricular hypertrophy in HO-1 -/- mice
10 than in HO+/+ mice, and (3) in HO-1 -/- (but not +/+ mice), massive cardiomyocyte death occurs with large organized thrombi attached to the infarct site. Although some cardiomyocyte death appears to be due to necrosis, apoptosis is a significant mechanism of cardiomyocyte
15 death. Hypoxia and elevated pulmonary arterial pressure increase cardiac production of reactive oxygen species, which play a significant role in myocardial death during ischemia/reperfusion.

Although myocardial infarction has been shown to
20 increase oxidative stress, a 2-3 fold increase in the nitration of protein tyrosine residues (which indicates the presence of the potent oxidant peroxynitrite) was detected in noninfarcted HO-1-deficient hearts exposed to 7 weeks of hypoxia. These data indicate that an increase
25 in oxidative stress precedes gross myocardial infarction.

Evidence of extensive lipid peroxidation in the zone of right ventricular infarction supports the conclusion that the absence of HO-1 in cardiomyocytes leads to an accumulation of reactive oxygen species that
30 causes cardiomyocyte death. Administration of HO-1 or overexpression of HO-1 protects against cardiomyocyte damage from hemodynamic stress or ischemia/reperfusion.
Adaptation of the cardiovascular system to hypoxia

The gene deletion studies described herein
35 indicate that HO-1 plays an important protective role in

- 13 -

vivo in the adaptation of the cardiovascular system to hypoxia. Right ventricles from HO-1 -/- mice were severely dilated and contained right ventricular infarcts with mural thrombi.

- 5 Humans and animals respond to hypoxia by exhibiting pulmonary vascular remodeling, pulmonary hypertension, and hypertrophy of the right ventricle. The data described herein were obtained using a mouse model of vascular injury which mimics the human response.
- 10 Hypoxia induces HO-1 expression in the lung, and CO generated by hypoxic VSMCs inhibits proliferation of these cells.

The data described herein indicate that the absence of HO-1 results in a maladaptive response in

15 cardiomyocytes exposed to hypoxia-induced pulmonary hypertension. In the absence of HO-1, VSMC are more sensitive to oxidative stress and have a maladaptive response to pressure overload. HO-1 has a protective effect on cardiomyocytes and VSMC subjected to stress

20 such as pressure-induced injury and secondary oxidative damage.

Therapeutic administration of HO

- In the absence of HO-1, cardiomyocytes undergo apoptotic cell death when subjected to stress such as
- 25 pressure overload or exposure to reactive oxygen species and that death can be inhibited by contacting the cells with HO. For example, HO-1, HO-2, or HO-3 protein or polypeptide (or DNA encoding HO-1, HO-2, or HO-3) is administered locally to heart tissue affected by hypoxic
- 30 conditions. One means for accomplishing local delivery is providing an HO or DNA encoding and HO on a surface of a vascular catheter, e.g., a balloon catheter coated with an antioxidant, which contacts the wall of the blood vessel to deliver therapeutic compositions at the site of

- 14 -

contact. Drug delivery catheters can also be used to administer solutions of therapeutic compositions.

HO-1 is therapeutically overexpressed (e.g., by administering an inducing agent to increase expression
5 from the endogenous gene) or by administering DNA (alone or in a plasmid) encoding an HO such as HO-1 or HO-2 (or an active fragment thereof, i.e., a fragment has the activity of inhibiting cardiomyocyte death). Inducing agents that stimulate HO-1 expression in cells include
10 hemin, hemoglobin, and heavy metals, e.g., SnCl_2 or NiCl_2 . For example, 250 mmol/kg of body weight of SnCl_2 or NiCl_2 is administered subcutaneously or 15 mg/kg of body weight of hemin is administered intraperitoneally to laboratory animals. Doses for human patients are determined and
15 optimized using standard methods.

Tables 1 and 3 show human HO-1 and HO-2 cDNA, respectively, in which the polypeptide-encoding nucleotides are designated in bold type and the termination codon is underlined. Tables 2 and 4 show the
20 amino acid sequences of human HO-1 and HO-2, respectively. Tables 5 and 6 show the nucleotide and amino acid sequence of rat HO-3.

- 15 -

TABLE 1: Human HO-1 cDNA

1 tcaacgcctg cctccccctcg agcgtcctca gcgcagccgc
cgccccgcgga gccagcacga
61 acgagcccag caccggccgg atggagcgtc cgcaaccoga
5 cagcatgccc caggatttgt
121 cagaggccct gaaggaggcc accaaggagg tgcacacca
ggcagagaat gctgagttca
181 tgaggaactt tcagaagggc caggtgaccc gagacggctt
caagctggtg atggcctccc
10 241 tgtaccacat ctatgtggcc ctggaggagg agattgagcg
caacaaggag agcccagtct
301 tcgccccctgt ctacttccca gaagagctgc accgcaaggc
tgccctggag caggacctgg
361 ccttctggta cgggccccgc tggcaggagg tcatccccta
15 cacaccagcc atgcagcgt
421 atgtgaagcg gctccacgag gtggggcgca cagagcccga
gctgctggtg gccacgcct
481 acaccgccta cctgggtgac ctgtctgggg gccaggtgct
caaaaagatt gccagaaag
20 541 ccctggacct gccagctct ggcgagggcc tggccttctt
caccttcccc aacattgcca
601 gtgccaccaa gttcaagcag ctctaccgct cccgcatgaa
ctccctggag atgactcccg
661 cagtcaggca gagggtgata gaagaggcca agactgcgtt
25 cctgctcaac atccagctct
721 ttgaggagtt gcaggagctg ctgacccatg acaccaagga
ccagagcccc tcacgggcac
781 cagggtctcg ccagcgggccc agcaacaaag tgcaagattc
tgcccccggtg gagactccca
30 841 gaggggaagcc cccactcaac acccgctccc aggctccgct
tctccgatgg gtccttacac
901 tcagctttct ggtggcgaca gttgctgtag ggctttatgc
catgtgaatg caggcatgct

- 16 -

961 ggctcccagg gccatgaact ttgtccggtg gaaggccttc
tttctagaga gggaattctc
1021 ttggctggct tccttaccgt gggcactgaa ggctttcagg
gcctccagcc ctctcactgt
5 1081 gtccctctct ctggaaagga ggaaggagcc tatggcatct
tccccaacga aaagcacatc
1141 caggcaatgg cctaaacttc agagggggcg aaggggtcag
ccctgccctt cagcatcctc
1201 agttcctgca gcagagcctg gaagacaccc taatgtggca
10 gctgtctcaa acctccaaaa
1261 gccctgagtt tcaagtatcc ttgttgacac ggccatgacc
actttccccc tgggccatgg
1321 caatttttac acaaacctga aaagatgttg tgtcttgtgt
ttttgtctta tttttgttgg
15 1381 agccactctg ttcttggtc agcctcaaat gcagtatctt
tggtgtgttc tgttgttttt
1441 atagcagggt tggggtggtt tttgagccat gcgtgggtgg
ggagggaggt gtttaacggc
1501 actgtggcct tgggtctaact tttgtgtgaa ataataaaca
20 acattgtctg
(SEQ ID NO:1)

Table 2: Human HO-1 amino acid sequence

MERPQPDSMP QDLSEALKEA TKEVHTQAEN AEFMRNFQKG QVTRDGFKL
MASLYHIYVA
25 LEEEIERNKE SPVFAPVYFP EELHRKAALE QDLAFWYGPR WQEVIPYTPA
MORYVKRLHE
VGRTEPELLV AHAYTRYLGD LSGGQVLKKI AQKALDLPSS GEGLAFFTFP
NIASATKFKQ
LYRSRMNSLE MTPAVRQRVI EEAKTAFLLN IQLFEELQEL LTHDTKDQSP
30 SRAPGLRQRA
SNKVQDSAPV ETPRGKPPLN TRSQAPLLRW VLTLNFLVAT VAVGLYAM (SEQ
ID NO:2)

- 17 -

Table 3: Human HO-2 cDNA

1 gggctgactg gaggctggcg gacaggcgac agacctgcgg
caggaccaga ggagcgagac
61 gagcaagaac cacacccagc agcaatgtca gcggaagtgg
5 aaacctcaga gggggtagac
121 gagtcagaaa aaaagaactc tggggcccta gaaaaggaga
accaaataag aatggctgac
181 ctctcagagc tcctgaagga agggaccaag gaagcacacg
accgggcaga aaacacccag
10 241 tttgtcaagg acttcttgaa aggcaacatt aagaaggagc
tgtttaagct ggccaccacg
301 gcactttact tcacatactc agccctcgag gaggaatatg
agcgcaacaa ggaccatcca
361 gcctttgccc ctttgtactt ccccatggag ctgcaccgga
15 aggaggcgct gaccaaggac
421 atggagtatt tctttggtga aaactgggag gagcaggtgc
agtgccccaa ggctgcccag
481 aagtacgtgg agcggatcca ctacataggg cagaacgagc
cggagctact ggtggcccat
20 541 gcatacacc gctacatggg ggatctctcg gggggccagg
tgctgaagaa ggtggcccag
601 cgagcactga aactccccag cacaggggaa gggacccagt
tctacctgtt tgagaatgtg
661 gacaatgccc agcagttcaa gcagctctac cgggccagga
25 tgaacgccct ggacctgaac
721 atgaagacca aagagaggat cgtggaggcc aacaaggctt
ttgagtataa catgcagata
781 ttcaatgaac tggaccaggc cggctccaca ctggccagag
agaccttgga ggatgggttc
30 841 cctgtacacg atgggaaagg agacatgcgt aaatgccctt
tctacgctgc tgaacaagac
901 aaagggtgg agggcagcct gtcccttccg acaagctatg
ctgtgctgag gaagcccagc

- 18 -

961 ctccagttca tcctggccgc tgggtgtggcc ctagctgctg
gactcttggc ctggtactac
1021 atgtgaagca cccatcatgc cacaccggta ccctcctccc
gactgaccac tggcctaccc
5 1081 ctttctccag ccttgactaa actaccacct caggtgactt
tttaaaaaat gctgggttta
1141 agaaaggcaa ccaataaaaag agatgctaga gcctcgtctg
acagcatcct ctctatgggc
1201 catattccgc actgggcaca ggccgtcacc ctgggagcag
10 tcggcacagt gcagcaagcc
1261 tggccccga cccagctcta ctccaggctt ccacacttct
gggccctagg ctgcttcgg
1321 tagtccctgt ttttgcagta catgggtgac tatctccct
gttggaggtg agtggcctgt
15 1381 aagtccaagc tgtgcgaggg ggcccttgctg gatgctgctg
tacaacttct gggcctctct
1441 tggaccctgg gagtgagggt ggggtgtgggt ggaagcctca
gaggccttgg gagctcatcc
1501 ctctcaccca gaatccctct aacccttggg tgcggtttgc
20 tcagccccag cttatctcct
1561 cctccgcctg tgtaaatgct ccagcactca ataaagtggg
ctttgcaagc taataaaaaa
1621 aaaaaaa (SEQ ID NO:3)

- 19 -

Table 4: Human HO-2 amino acid sequence

MSAEVETSEGVDESEKKNSGALEKENQMRMADLSELLKEGTKEAHDRAENTQFVKDF
LKGNIKKELFKLATTALYFTYSALEEEMERNKDHFAFAPLYFPMELHRKEALTKDME
YFFGENWEEQVQCPKAAQKYVERIHYIGQNEPELLVAHAYTRYMGDLSGGQVLKKVA
5 QRALKLPSTGEGTQFYLFENVDNAQQFKQLYRARMNALDLNMKTKERIVEANKAFEY
NMQIFNELDQAGSTLARETLEDGFPVHDGKGDMRKCPFYAAEQDKGLEGSLSLPTSY
AVLRKPSLQFILAAGVALAAGLLAWYYM (SEQ ID NO:4)

Table 5: Rat HO-3 nucleotide sequence

1 tttcagggat ttttgcgatt cctctctgta gacttctact
10 tgttctctaa gggagttctt
61 catgtctttc ttgaagtcac ccagcatcat gatcaaatat
gattttgaaa ctagatcttg
121 cttttctggg gtgtttggat attccatggt tgttttggtg
ggagaattgg gctccgatga
15 181 tggcatgtag tcttggtttc tgttgcttgg tttcctgcgc
ttgcctctcg ccatcagatt
241 atctctagtg ttactttggt ctgctatttc tgacagtggc
tagactgtcc tataagcctg
301 tgtgtcagga gtgctgtaga ctttttttcc tctctttcag
20 tcagttatgg gacagagtgt
361 tctgcttttg ggcgtgtagt ttttcctctc tacaggtctt
cagctgttcc tgtgggcctg
421 tgtcttgagt tcaccaggca gctttcttgc agcagaaaat
ttggtcatac ctgtgatcct
25 481 gaggetcaag ttcgctcgtg ggggtgctgtc caggggctct
ctgcagcggg cacaaccagg
541 aagacctgtg cggccccttc cggagcttca gtgcaccagg
gttccagatg gcctttggcg
601 ttttcctctg gcgtccgaga tgtatgtaca gagagcagtc
30 tcttctgggt tcccaggctt
661 gtctgcctct ctgaagggtc agctctccct cccacgggat
ttgggtgcag agaactgttt

- 20 -

721 atccggtctg tttctttcag gttccggtgg tgtctcaggc
aggtgtcgtt cctgcgccct
781 ccccatggg accagaggcc ttatacagtt tcctcttggg
ccagggatgt gggcaggggt
5 841 gagcagtgtt ggtggtctct tccgtctgca gcctcaggag
tgccacctga ccaggcgggt
901 gggctctctct ctgagaattt cattttttaa tcattcatta
aaatgtcatg acttgatgtc
961 ctgctgtccg tctcacgccc tcagctgtaa cagtgccgag
10 ggagtcactg aagaagagac
1021 tgaatgacca gagtatgggc agcacagaca actcaacaaa
aatgtcttca gaggtggaga
1081 ctgcggaggc cgtagatgag tcagagaaga actctatggc
atcagagaag gaaaaccatt
15 1141 caaaatagc agacttttct gatcttctga aggaaggagc
aaaggaagca gatgaccggg
1201 cagaaaatac ccagtttgtc aaagacttct tgaaaggaaa
cattaagaag gagctattta
1261 agctggccac cactgcactt tcatactcag cccctgagga
20 ggaaatggat tcaactgacca
1321 aggacatgga gtacttcttt ggtgaaaact gggaggaaaa
agtgaagtgc tctgaagctg
1381 cccagacgta tgtggatcag attcactatg tagggcaaaa
tgagccagag catctgggtg
25 1441 cccatactta ctctacttac atggggggaa acctttcagg
ggaccaggta ctgaagaagg
1501 agaccagcc ggtccccttc actagggaag ggactcagtt
ctacctgttt gagcatgtag
1561 acaatgctaa gcaattcaag ctattctact gcgctagatt
30 gaatgccttg gacctgaatt
1621 tgaagaccaa agagaggatt gtggaggaag ccaccaaaagc
ctttgaatat aatatgcaga
1681 tattcagtga actggaccag gcaggctcca taccagtaag
agaaacccta aagaatgggc

- 21 -

1741 tctcaatact tgatgggaag ggaggtgtat gcaaagtcc
 ctttaatgct gctcagccag
 1801 acaaaggtac cctgggaggc agcaactgcc ctttccagat
 gtccatggcc ttgctgagga
 5 1861 agcctaactt gcagctcatt ctagttgcca gtatggcctt
 ggtagctgga cttttagcct
 1921 ggtactacat gtgaagggcc tgtcaagttg tttgcatcct
 atctcaacat cctaccactt
 1981 gttccttccc cacctccacc tctgcctaga actaccacct
 10 caggtgacat ttttaatggt
 2041 gggtttgaga aaatgagcaa ccaataaaaag acagacccta
 gaaaaaagtc atgacttaag
 2101 tggcacgggg acacctaaag tcacactttg tgcttcagac
 atactttctt tctctatttc
 15 2161 aacactgaat tcgggaagta acctactact attaataata
 aatgctacac aatgcataat
 2221 aaaaa (SEQ ID NO:5)

Table 6: Rat HO-3 amino acid sequence

MSSEVETAEAVDESEKNSMASEKENHSKIADFSDLLKEGTKEADDRAENTQFVKDFL
 20 KGNIKKELFKLATTALSYSAPEEEMDSLTKDMEYFFGENWEEKVKCSEAAQTYVDQI
 HYVGQNEPEHLVAHTYSTYMGGNLSGDQVLKKETQVPVFTREGTQFYLFHVNDNAKQ
 FKLIFYCARLNALDLNLKTKERIVEEATKAFEYNMQIFSELDQAGSIPVRETLKNGLS
 ILDGKGGVCKCPFNAAPDKGTLGGSNCPFQMSMALLRKPNLQLILVASMALVAGLL
 AWYYM (SEQ ID NO:6)

25 An HO preferably has an amino acid sequence that
 is at least 85% identical (preferably at least 90%, more
 preferably at least 95%, more preferably at least 98%,
 most preferably at least 100% identical) to the amino
 acid sequence of SEQ ID NO: 2, 4, or 6. DNA encoding an
 30 HO preferably has nucleotide sequence that is at least
 50% identical (preferably at least 75%, more preferably
 at least 85%, more preferably at least 95%, most

- 22 -

preferably at least 100% identical) to the nucleotide sequence of the coding region of SEQ ID NO:1, 3, or 5.

The per cent identity of nucleotide and amino acid sequences is determined using the Sequence Analysis

5 Software Package developed by the Genetics Computer Group (University of Wisconsin Biotechnology Center, Madison, WI), employing the default parameters thereof.

To be clinically beneficial, expression of HO from an endogenous gene or expression of recombinant HO from
10 exogenous DNA need not be long term. For example, to inhibit cardiomyocyte death associated with a myocardial infarction or other acute condition which results in oxidative stress to the tissue, the most critical period of treatment is the first three months after injury.

15 Thus, gene therapy to express recombinant HO for even a period of days or weeks after administration of the HO-encoding DNA (which administration is prior to or soon after an injury) minimizes cell death, inhibits VSMC proliferation, and therefore confers a clinical benefit.

20 For local administration of DNA to cardiovascular tissue, standard gene therapy vectors used. Such vectors include viral vectors, including those derived from replication-defective hepatitis viruses (e.g., HBV and HCV), retroviruses (see, e.g., WO 89/07136; Rosenberg et
25 al., 1990, N. Eng. J. Med. 323(9):570-578), adenovirus (see, e.g., Morsey et al., 1993, J. Cell. Biochem., Supp. 17E,), adeno-associated virus (Kotin et al., 1990, Proc. Natl. Acad. Sci. USA 87:2211-2215,), replication
30 defective herpes simplex viruses (HSV; Lu et al., 1992, Abstract, page 66, Abstracts of the Meeting on Gene Therapy, Sept. 22-26, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York), and any modified versions of these vectors. The invention may utilize any other delivery system which accomplishes *in vivo* transfer of
35 nucleic acids into eukaryotic cells. For example, the

- 23 -

nucleic acids may be packaged into liposomes, e.g., cationic liposomes (Lipofectin), receptor-mediated delivery systems, non-viral nucleic acid-based vectors, erythrocyte ghosts, or microspheres (e.g.,

5 microparticles; see, e.g., U.S. Patent No. 4,789,734; U.S. Patent No. 4,925,673; U.S. Patent No. 3,625,214; Gregoriadis, 1979, *Drug Carriers in Biology and Medicine*, pp. 287-341 (Academic Press,)). Naked DNA may also be administered. Alternatively, a plasmid which directs

10 cardiospecific expression (e.g., a plasmid containing a myosin heavy chain (α MHC) promoter; Fig. 6) of an HO-encoding sequence can be used for gene therapy. Expression of an HO (encoded, e.g., by the coding sequences of SEQ ID NO:1, 3, or 5) from such a

15 constitutive promoter is useful to inhibit cardiomyocyte death *in vivo*. Nucleic acids which hybridize at high stringency to the coding sequences of SEQ ID NO:1, 3, or 5 and which encode a polypeptide which has a biological activity of an HO polypeptide (e.g., inhibition of

20 cardiomyocyte death) are also used for gene therapy for vascular injury. To determine whether a nucleic acid hybridizes to a reference nucleic acid at a given stringency, hybridization is carried out using standard techniques, such as those described in Ausubel *et al.*

25 (*Current Protocols in Molecular Biology*, John Wiley & Sons, 1989). "High stringency" refers to nucleic acid hybridization and wash conditions characterized by high temperature and low salt concentration, e.g., wash conditions of 65°C at a salt concentration of

30 approximately 0.1 X SSC. "Low" to "moderate" stringency refers to DNA hybridization and wash conditions characterized by low temperature and high salt concentration, e.g., wash conditions of less than 60°C at a salt concentration of at least 1.0 X SSC. For example,

35 high stringency conditions may include hybridization at

- 24 -

about 42°C, and about 50% formamide; a first wash at about 65°C, about 2X SSC, and 1% SDS; followed by a second wash at about 65°C and about 0.1% x SSC. Lower stringency conditions suitable for detecting DNA

- 5 sequences having about 50% sequence identity to an CHF-1 gene are detected by, for example, hybridization at about 42°C in the absence of formamide; a first wash at about 42°C, about 6X SSC, and about 1% SDS; and a second wash at about 50°C, about 6X SSC, and about 1% SDS. To
- 10 determine whether a polypeptide encoded by a hybridizing nucleic acid has a biological activity of an HO polypeptide, the polypeptide is evaluated using any of the functional assays to measure HO activity described herein, e.g., measuring VSMC proliferation or
- 15 cardiomyocyte death.

For gene therapy of cardiovascular tissue, fusigenic viral liposome delivery systems known in the art (e.g., hemagglutinating virus of Japan (HVJ) liposomes or Sendai virus-liposomes) are useful for

20 efficiency of plasmid DNA transfer (Dzau et al., 1996, Proc. Natl. Acad. Sci. U.S.A. 93:11421-11425). Using HVJ-liposomes, genes are expressed from plasmid DNA delivered to target tissues *in vivo* for extended periods of time (e.g., greater than two weeks for heart and

25 arterial tissue and up to several months in other tissues).

DNA for gene therapy can be administered to patients parenterally, e.g., intravenously, subcutaneously, intramuscularly, and intraperitoneally.

30 Sustained release administration such as depot injections or erodible implants, e.g., vascular stents coated with DNA encoding an HO, may also be used. The compounds may also be directly applied during surgery, e.g., bypass surgery, or during angioplasty, e.g., an angioplasty

35 catheter may be coated with DNA encoding an HO. The DNA

- 25 -

is then deposited at the site of angioplasty. DNA or an inducing agent is administered in a pharmaceutically acceptable carrier, i.e., a biologically compatible vehicle, which is suitable for administration to an animal e.g., physiological saline. A therapeutically effective amount is an amount which is capable of producing a medically desirable result, e.g., expression of HO, in a treated animal. Such an amount can be determined by one of ordinary skill in the art. As is well known in the medical arts, dosage for any given patient depends upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, severity of arteriosclerosis or vascular injury, and other drugs being administered concurrently. Dosages will vary, but a preferred dosage for intravenous administration of DNA is approximately 10^6 to 10^{22} copies of the DNA molecule.

HO-based therapy for cardiovascular disorders depends on when (in the course of a vascular injury) the patient is encountered. HO-1 -/- VSMC initially proliferated at a faster rate compared to wild type VSMC within days after an insult. Increasing local HO-1 levels at this stage inhibits VSMC growth and confers a clinical benefit. For example, if the patient is encountered at an early stage (e.g., within one week of cardiovascular stress or injury), the patient is treated by augmenting the local level of HO-1 (e.g., by administering an HO polypeptide, by increasing expression of endogenous HO, or by standard gene therapy techniques described above to produce recombinant HO *in vivo*) to inhibit the growth of VSMC and decrease the size of a myocardial infarct. In contrast, if the patient is encountered at a later stage (e.g., several weeks, 1 month, 2 months, and up to 3 months after an injury), the

- 26 -

patient is treated by inhibiting HO expression to decrease formation of a stenotic lesion, e.g., by antisense therapy, as described below.

Organ and tissue preservation

5 Concerns about irreversible ischemic tissue damage arise when a donor organ, e.g., a heart, is removed from the donor and stored for more than about 5-6 hours before transplantation into the recipient. *Ex vivo* treatment of a donor organ to reduce tissue damage by inhibiting death
10 of cardiomyocytes is carried out by immersing the organ in a solution containing an inducing agent, an HO, e.g., HO-1 or HO-2, or a nucleic acid encoding an HO prior to transplantation. By "ex vivo treatment" is meant treatment that takes place outside of the body. For
15 example, ex vivo treatment is administered to an organ (or a tissue fragment or dissociated cells) that has been removed from a mammal and which will be returned to the same or a different mammal (at the same or different anatomical site) after the treatment. For example,
20 effective DNA delivery to cells in solid organs achieved by contacting the organ with a combination of DNA, liposomes and transferrin during the cold ischemic time prior to transplantation (see, e.g., Hein et al. 1998, Eur. J. Cardiothorac. Surg. 13:460-466). An organ may
25 also be perfused or electroporated with solution containing HO-encoding DNA. Vectors and delivery systems described above for gene therapy applications are suitable for organ preservation and cell preservation *in vitro*.

30 Inhibition of restenosis

VSMC proliferation contributes to graft stenosis and restenosis following vascular injury such as that resulting from coronary angioplasty and coronary bypass surgery. Patients with restenosis have a significantly

- 27 -

poorer clinical outcome compared to patients without restenosis.

Using a mouse model of vascular graft stenosis in which the stenosis develops rapidly and closely mimics the development of vascular graft stenosis in humans, the effect of HO-1 on VSMC proliferation was examined. A patch of jugular vein was grafted onto a carotid artery in normal and HO-1 deficient mice to create composite vessels that mimic vein grafts used for bypass surgery (Fig. 4). The vein patch is subject to increased pressure which leads to an increase in local VSMC proliferation and occlusion of the blood vessel (Figs. 5A-B). In wild type mice, there was robust formation of a neointima (characterized by proliferating VSMC) in the vein graft. In contrast, tissue sections of the neointima of HO-1 -/- mice revealed a necrotic mass. The HO-1 -/- neointima was a complex lesion characterized by mostly acellular material, indicating death of VSMC. HO-1 -/- VSMC are more susceptible to H₂O₂-induced death compared to VSMC isolated from wild type mice (Fig. 8). These data indicate that HO-1 is required for VSMC proliferation at later stages post-injury and that local inhibition of HO-1 expression, e.g., by antisense therapy, is useful to inhibit graft stenosis or restenosis in patients affected or who at risk of developing such conditions.

The data described herein indicate that (1) in response to increased pressure, VSMC proliferate in the neointima of the venous patch in HO-1 +/+ mice, and (2) in contrast, massive cell death occurs in the neointima of the venous patch in HO-1 -/- mice.

Patients undergoing invasive vascular procedures, e.g., balloon angioplasty, are at risk for developing undesired vascular stenosis or restenosis. Angioplasty, used to treat arteriosclerosis, involves the insertion of

- 28 -

catheters, e.g., balloon catheters, through an occluded region of a blood vessel in order to expand it. However, the aftermath of angioplasty may be problematic.

Restenosis, or closing of the vessel, can occur as a

5 consequence of injury, e.g., mechanical abrasion

associated with the angioplasty treatment. This

restenosis is caused by proliferation of smooth muscle

cells stimulated by vascular injury. Other anatomical

disruptions or mechanical disturbances of a blood

10 vessel, e.g., laser angioplasty, coronary artery surgery,

atherectomy, coronary artery stents, and coronary bypass

surgery, may also cause vascular injury and subsequent

proliferation of smooth muscle cells.

Therapeutic approaches, such as antisense therapy

15 or ribozyme therapy are used to inhibit HO expression,

and as a result, VSMC proliferation that leads to

neointimal thickening. The antisense strand (either RNA

or DNA) is directly introduced into the cells in a form

that is capable of binding to the mRNA transcripts.

20 Alternatively, a vector-containing sequence which, which

once within the target cells is transcribed into the

appropriate antisense mRNA, may be administered. Nucleic

acids complementary to all or part of the HO cDNA (SEQ ID

NO: 1, 3, or 5) may be used in methods of antisense

25 treatment to inhibit expression of HO. Antisense

treatment is carried out by administering to a mammal,

such as a human, DNA containing a promoter, e.g., a

cardiospecific promoter, operably linked to a DNA

sequence (an antisense template), which is transcribed

30 into an antisense RNA. By "operably linked" is meant

that a coding sequence and a regulatory sequence(s)

(i.e., a promoter) are connected in such a way as to

permit gene expression when the appropriate molecules

(e.g., transcriptional activator proteins) are bound to

35 the regulatory sequence(s). Alternatively, as mentioned

- 29 -

above, antisense oligonucleotides may be introduced directly into vascular cells. The antisense oligonucleotide may be a short nucleotide sequence (generally at least 10, preferably at least 14, more preferably at least 20 (e.g., at least 30), and up to 100 or more nucleotides) formulated to be complementary to a portion, e.g., the coding sequence, or all of HO mRNA. Oligonucleotides complementary to various portions of HO-1 or HO-2 mRNA can readily be tested *in vitro* for their ability to decrease production of HO in cells, using standard methods. Sequences which decrease production of HO message in *in vitro* cell-based or cell-free assays can then be tested *in vivo* in rats or mice to determine whether HO expression (or VSMC proliferation) is decreased.

Ribozyme therapy can also be used to inhibit gene expression. Ribozymes bind to specific mRNA and then cut it at a predetermined cleavage point, thereby destroying the transcript. These RNA molecules may be used to inhibit expression of a gene encoding a protein involved in the formation of vein graft stenosis according to methods known in the art (Sullivan et al., 1994, J. Invest. Derm. 103:85S-89S; Czubayko et al., 1994, J. Biol. Chem. 269:21358-21363; Mahieu et al, 1994, Blood 84:3758-65; Kobayashi et al. 1994, Cancer Res. 54:1271-1275).

Standard methods of administering antisense therapy have been described (see, e.g., Melani et al., 1991, Cancer Res. 51:2897-2901). Antisense nucleic acids which hybridize to HO-encoding mRNA can decrease or inhibit production of HO by associating with the normally single-stranded mRNA transcript, thereby interfering with translation and thus, expression of HO. Such nucleic acids are introduced into target cells by standard vectors and/or gene delivery systems such as those

- 30 -

described above for gene therapy. Suitable gene delivery systems may include liposomes, receptor-mediated delivery systems, naked DNA, and viral vectors such as herpes viruses, retroviruses, adenoviruses and adeno-associated viruses, among others. For example, antisense oligodesoxynucleotides, e.g., oligonucleotides which have been modified to phosphorothioates or phosphoramidates, withstand degradation after delivery and have been successfully used to inhibit gene expression in a model of reperfusion injury (see, e.g., Haller et al., 1998, *Kidney Int.* 53:1550-1558). Pharmaceutically acceptable carriers are biologically compatible vehicles which are suitable for administration to an animal: e.g., physiological saline. A therapeutically effective amount of a compound is an amount which is capable of producing a medically desirable result in a treated animal, e.g., inhibition of expression of HO-1 or a decrease in VSMC proliferation.

Compositions that inhibit HO activity, e.g., its role in the promotion of VSMC proliferation, are also administered to inhibit VSMC-mediated stenosis or restenosis. For example, metalloporphyrins, e.g., zinc protoporphyrin IX (ZnPP), zinc mesoporphyrin IX (ZnMP), tin protoporphyrin IX (SnPP), tin mesoporphyrin IX (SnMP), zinc deuteroporphyrin IX 2,4 bis glycol (ZnDPBG), chromium protoporphyrin (CrPP), cobalt protoporphyrin (CoPP), and manganese metalloporphyrin (MnPP) are administered to mammals at $\mu\text{mol/kg}$ doses to inhibit HO activity. SnPP has safely been administered to human infants at doses of $0.5 \mu\text{mol/kg}$ to $100 \mu\text{mol/kg}$ of body weight. HO-inhibitory doses for local administration are determined using methods known in the art.

Parenteral administration, such as intravenous, subcutaneous, intramuscular, and intraperitoneal delivery routes, may be used to deliver the compounds that inhibit

- 31 -

HO activity or expression, with local vascular administration being the preferred route. Dosages for any one patient depends upon many factors, including the patient's size, body surface area, age, the particular
5 compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently. For antisense therapy, a preferred dosage for administration of nucleic acids is from approximately 10^6 to 10^{22} copies of the nucleic acid
10 molecule. As described above, local administration to a site of vascular injury or to cardiac tissue is accomplished using a catheter or indwelling vascular stent.

Other embodiments are within the following claims.

15 What is claimed is:

- 32 -

1. A method of inhibiting cardiomyocyte death in a mammal comprising locally administering to the myocardium of said mammal a heme oxygenase (HO).

2. The method of claim 1, wherein said mammal has suffered a myocardial infarction.

3. The method of claim 1, wherein said mammal has myocarditis.

4. The method of claim 1, wherein said HO is heme oxygenase-1 (HO-1).

5. The method of claim 1, wherein said HO is heme oxygenase-2 (HO-2).

6. A method of inhibiting cardiomyocyte death in a mammal comprising locally administering to the myocardium of said mammal a DNA encoding a HO.

7. The method of claim 6, wherein said HO is HO-1.

8. The method of claim 6, wherein said HO is HO-2 or HO-3.

9. A method of inhibiting cardiomyocyte death *in vitro*, comprising contacting cardiomyocytes with an HO.

10. A method of inhibiting cardiomyocyte death *in vitro*, comprising contacting cardiomyocytes with DNA encoding an HO.

- 33 -

11. The method of claim 10, wherein said HO is HO-1.

12. The method of claim 10, wherein said HO is HO-2.

5 13. A method of preserving isolated myocardial tissue comprising perfusing said tissue with a solution comprising an HO or a DNA encoding an HO.

14. A method of inhibiting vascular restenosis in a mammal comprising locally administering to the site of
10 a vascular injury a compound which inhibits expression of HO-1.

15. The method of claim 14, wherein said compound inhibits HO-1 transcription in a vascular cell of said mammal.

15 16. The method of claim 15, wherein said vascular cell is an aortic smooth muscle cell.

17. The method of claim 14, wherein said mammal is a human.

18. The method of claim 14, wherein said compound
20 inhibits translation of HO-1 mRNA in a vascular cell of said mammal.

19. The method of claim 18, wherein said compound consists of a single stranded nucleic acid complementary to at least a portion of said HO-1 mRNA.

- 34 -

20. A method of inhibiting vascular restenosis in a mammal comprising identifying a mammal having vascular restenosis or at risk of developing vascular restenosis and administering to said mammal a compound which
5 inhibits expression of HO-1.

21. The method of claim 14, wherein said compound is administered to said mammal at least one month after a vascular injury.

22. The method of claim 14, wherein said compound
10 is administered to said mammal at least two months after a vascular injury.

23. The method of claim 14, wherein said compound is administered to said mammal at least three months after a vascular injury.

15 24. A method of inhibiting vascular smooth muscle cell proliferation in a mammal comprising administering to an injured vascular tissue of said mammal a HO, wherein said HO is administered within 24 hours after a vascular injury.

20 25. The method of claim 24, wherein said HO is administered to said mammal for up to one week after a vascular injury.







FIG. 1


2/8




3/8

**FIG. 4**


4/8

FIG. 5A**FIG. 5B**


5/8

d-MHC α
promoter**FIG. 6**



6/8

**FIG. 7**

7/8

**FIG. 8**

8/8



SEQUENCE LISTING

<110> The President and Fellows of Harvard College

<120> INHIBITING CARDIOMYOCYTE DEATH

<130> 00246/235WO2

<150> US 60/121,946

<151> 1999-02-25

<150> US 60/098,377

<151> 1998-08-28

<160> 6

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 1550

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (81)...(944)

<400> 1

tcaacgcctg cctccctcg agcgtctca ggcagccgc cgcccgcgga gccagcacga	60
acgagcccag caccggccgg atg gag cgt ccg caa ccc gac agc atg ccc cag	113
Met Glu Arg Pro Gln Pro Asp Ser Met Pro Gln	
1 5 10	
gat ttg tca gag gcc ctg aag gag gcc acc aag gag gtg cac acc cag	161
Asp Leu Ser Glu Ala Leu Lys Glu Ala Thr Lys Glu Val His Thr Gln	
15 20 25	
gca gag aat gct gag ttc atg agg aac ttt cag aag ggc cag gtg acc	209
Ala Glu Asn Ala Glu Phe Met Arg Asn Phe Gln Lys Gly Gln Val Thr	
30 35 40	
cga gac ggc ttc aag ctg gtg atg gcc tcc ctg tac cac atc tat gtg	257
Arg Asp Gly Phe Lys Leu Val Met Ala Ser Leu Tyr His Ile Tyr Val	
45 50 55	
gcc ctg gag gag gag att gag cgc aac aag gag agc cca gtc ttc gcc	305
Ala Leu Glu Glu Glu Ile Glu Arg Asn Lys Glu Ser Pro Val Phe Ala	
60 65 70 75	
cct gtc tac ttc cca gaa gag ctg cac cgc aag gct gcc ctg gag cag	353
Pro Val Tyr Phe Pro Glu Glu Leu His Arg Lys Ala Ala Leu Glu Gln	
80 85 90	
gac ctg gcc ttc tgg tac ggg ccc cgc tgg cag gag gtc atc ccc tac	401
Asp Leu Ala Phe Trp Tyr Gly Pro Arg Trp Gln Glu Val Ile Pro Tyr	
95 100 105	
aca cca gcc atg cag cgc tat gtg aag cgg ctc cac gag gtg ggg cgc	449
Thr Pro Ala Met Gln Arg Tyr Val Lys Arg Leu His Glu Val Gly Arg	
110 115 120	
aca gag ccc gag ctg ctg gtg gcc cac gcc tac acc cgc tac ctg ggt	497
Thr Glu Pro Glu Leu Leu Val Ala His Ala Tyr Thr Arg Tyr Leu Gly	
125 130 135	

gac Asp 140	ctg Leu	tct Ser	ggg Gly	ggc Gly	cag Gln 145	gtg Val	ctc Leu	aaa Lys	aag Lys	att Ile 150	gcc Ala	cag Gln	aaa Lys	gcc Ala	ctg Leu 155	545	
gac Asp	ctg Leu	ccc Pro	agc Ser	tct Ser 160	ggc Gly	gag Glu	ggc Gly	ctg Leu	gcc Ala 165	ttc Phe	ttc Phe	acc Thr	ttc Phe	ccc Pro 170	aac Asn	593	
att Ile	gcc Ala	agt Ser	gcc Ala 175	acc Thr	aag Lys	ttc Phe	aag Lys	cag Gln 180	ctc Leu	tac Tyr	cgc Arg	tcc Ser	cgc Arg 185	atg Met	aac Asn	64	
tcc Ser	ctg Leu	gag Glu 190	atg Met	act Thr	ccc Pro	gca Ala	gtc Val 195	agg Arg	cag Gln	agg Arg	gtg Val	ata Ile 200	gaa Glu	gag Glu	gcc Ala	689	
aag Lys	act Thr 205	gcg Ala	ttc Phe	ctg Leu	ctc Leu	aac Asn 210	atc Ile	cag Gln	ctc Leu	ttt Phe	gag Glu 215	gag Glu	ttg Leu	cag Gln	gag Glu	737	
ctg Leu 220	ctg Leu	acc Thr	cat His	gac Asp	acc Thr 225	aag Lys	gac Asp	cag Gln	agc Ser	ccc Pro 230	tca Ser	cgg Arg	gca Ala	cca Pro	ggg Gly 235	785	
ctt Leu	cgc Arg	cag Gln	cgg Arg	gcc Ala 240	agc Ser	aac Asn	aaa Lys	gtg Val	caa Gln 245	gat Asp	tct Ser	gcc Ala	ccc Pro	gtg Val 250	gag Glu	833	
act Thr	ccc Pro	aga Arg	ggg Gly 255	aag Lys	ccc Pro	cca Pro	ctc Leu	aac Asn 260	acc Thr	cgc Arg	tcc Ser	cag Gln	gct Ala 265	ccg Pro	ctt Leu	881	
ctc Leu	cga Arg	tgg Trp 270	gtc Val	ctt Leu	aca Thr	ctc Leu	agc Ser 275	ttt Phe	ctg Leu	gtg Val	gcg Ala	aca Thr 280	gtt Val	gct Ala	gta Val	929	
ggg Gly 285	ctt Leu	tat Tyr	gcc Ala	atg Met	tgaatgcagg			catgctggct			cccagggcca			tgaactttgt			984
ccggtggaag			gccttctttc			tagagaggga			attctcttgg			ctggcttcc			taccgtgggc		1044
actgaaggct			ttcagggcct			ccagccctct			cactgtgtcc			ctctctctgg			aaaggaggaa		1104
ggagcctatg			gcatcttccc			caacgaaaag			cacatccagg			caatggccta			aacttcagag		1164
ggggcgaagg			ggtcagccct			gcccttcagc			atcctcagtt			cctgcagcag			agcctggaag		1224
acaccctaata			gtggcagctg			tctcaaacct			ccaaaagccc			tgagtttcaa			gtatccttgt		1284
tgacacggcc			atgaccactt			tccccgtggg			ccatggcaat			ttttacacaa			acctgaaaag		1344
atgttggtg			ttgtgttttt			gtcttatttt			tgttggagcc			actctgttcc			tggtcagcc		1404
tcaaatagcag			tattttttgt			gtgttctgtt			gtttttatag			cagggttggg			gtgggtttttg		1464
agccatgcgt			gggtggggag			ggaggtgttt			aacggcactg			tggccttggt			ctaacttttg		1524
tgtgaaataa			taaacaacat			tgtctg											

```
<210> 2
<211> 288
<212> PRT
<213> Homo sapiens
```

<400> 2															
Met	Glu	Arg	Pro	Gln	Pro	Asp	Ser	Met	Pro	Gln	Asp	Leu	Ser	Glu	Ala
1				5					10					15	
Leu	Lys	Glu	Ala	Thr	Lys	Glu	Val	His	Thr	Gln	Ala	Glu	Asn	Ala	Glu
			20					25					30		
Phe	Met	Arg	Asn	Phe	Gln	Lys	Gly	Gln	Val	Thr	Arg	Asp	Gly	Phe	Lys
		35					40					45			
Leu	Val	Met	Ala	Ser	Leu	Tyr	His	Ile	Tyr	Val	Ala	Leu	Glu	Glu	Glu
	50					55					60				
Ile	Glu	Arg	Asn	Lys	Glu	Ser	Pro	Val	Phe	Ala	Pro	Val	Tyr	Phe	Pro
65					70					75					80

Glu	Glu	Leu	His	Arg 85	Lys	Ala	Ala	Leu	Glu 90	Gln	Asp	Leu	Ala	Phe 95	Trp
Tyr	Gly	Pro	Arg	Trp	Gln	Glu	Val	Ile 105	Pro	Tyr	Thr	Pro	Ala	Met	Gln
Arg	Tyr	Val	Lys	Arg	Leu	His	Glu 120	Val	Gly	Arg	Thr	Glu 125	Pro	Glu	Leu
Leu	Val	Ala	His	Ala	Tyr	Thr 135	Arg	Tyr	Leu	Gly	Asp 140	Leu	Ser	Gly	Gly
Gln 145	Val	Leu	Lys	Lys	Ile 150	Ala	Gln	Lys	Ala	Leu 155	Asp	Leu	Pro	Ser	Ser 160
Gly	Glu	Gly	Leu	Ala 165	Phe	Phe	Thr	Phe	Pro 170	Asn	Ile	Ala	Ser	Ala 175	Thr
Lys	Phe	Lys	Gln 180	Leu	Tyr	Arg	Ser	Arg 185	Met	Asn	Ser	Leu	Glu 190	Met	Thr
Pro	Ala	Val	Arg	Gln	Arg	Val	Ile 200	Glu	Glu	Ala	Lys	Thr 205	Ala	Phe	Leu
Leu	Asn 210	Ile	Gln	Leu	Phe	Glu 215	Glu	Leu	Gln	Glu	Leu 220	Thr	His	Asp	
Thr 225	Lys	Asp	Gln	Ser	Pro 230	Ser	Arg	Ala	Pro	Gly 235	Leu	Arg	Gln	Arg	Ala 240
Ser	Asn	Lys	Val	Gln 245	Asp	Ser	Ala	Pro	Val 250	Glu	Thr	Pro	Arg	Gly 255	Lys
Pro	Pro	Leu	Asn 260	Thr	Arg	Ser	Gln	Ala 265	Pro	Leu	Leu	Arg	Trp 270	Val	Leu
Thr	Leu	Ser 275	Phe	Leu	Val	Ala 280	Thr	Val	Ala	Val	Gly	Leu 285	Tyr	Ala	Met

```
<210> 3
<211> 1627
<212> DNA
<213> Homo sapiens
```

```
<220>
<221> CDS
<222> (85)...(1023)
```

<400> 3																60
gggctgactg	gaggctggcg	gacaggcgac	agacctgcgg	caggaccaga	ggagcgagac											111
gagcaagaac	cacacccagc	agca	atg	tca	gcg	gaa	gtg	gaa	acc	tca	gag					
			Met	Ser	Ala	Glu	Val	Glu	Thr	Ser	Glu					
			1				5									
ggg	gta	gac	gag	tca	gaa	aaa	aag	aac	tct	ggg	gcc	cta	gaa	aag	gag	159
Gly	Val	Asp	Glu	Ser	Glu	Lys	Lys	Asn	Ser	Gly	Ala	Leu	Glu	Lys	Glu	
10					15					20					25	
aac	caa	atg	aga	atg	gct	gac	ctc	tca	gag	ctc	ctg	aag	gaa	ggg	acc	207
Asn	Gln	Met	Arg	Met	Ala	Asp	Leu	Ser	Glu	Leu	Leu	Lys	Glu	Gly	Thr	
				30					35					40		
aag	gaa	gca	cac	gac	cgg	gca	gaa	aac	acc	cag	ttt	gtc	aag	gac	ttc	255
Lys	Glu	Ala	His	Asp	Arg	Ala	Glu	Asn	Thr	Gln	Phe	Val	Lys	Asp	Phe	
			45					50					55			
ttg	aaa	ggc	aac	att	aag	aag	gag	ctg	ttt	aag	ctg	gcc	acc	acg	gca	303
Leu	Lys	Gly	Asn	Ile	Lys	Lys	Glu	Leu	Phe	Lys	Leu	Ala	Thr	Thr	Ala	
		60					65					70				
ctt	tac	ttc	aca	tac	tca	gcc	ctc	gag	gag	gaa	atg	gag	cgc	aac	aag	351
Leu	Tyr	Phe	Thr	Tyr	Ser	Ala	Leu	Glu	Glu	Glu	Met	Glu	Arg	Asn	Lys	
	75					80					85					
gac	cat	cca	gcc	ttt	gcc	cct	ttg	tac	ttc	ccc	atg	gag	ctg	cac	cgg	399
Asp	His	Pro	Ala	Phe	Ala	Pro	Leu	Tyr	Phe	Pro	Met	Glu	Leu	His	Arg	
90					95					100					105	

aag gag gcg ctg acc aag gac atg gag tat ttc ttt ggt gaa aac tgg	447
Lys Glu Ala Leu Thr 110 Lys Asp Met Glu Tyr 115 Phe Phe Gly Glu Asn Trp 120	
gag gag cag gtg cag tgc ccc aag gct gcc cag aag tac gtg gag cgg	495
Glu Glu Gln Val 125 Gln Cys Pro Lys Ala 130 Ala Gln Lys Tyr 135 Val Glu Arg	
atc cac tac ata ggg cag aac gag ccg gag cta ctg gtg gcc cat gca	543
Ile His Tyr 140 Ile Gly Gln Asn Glu Pro Glu Leu Leu Val Ala His Ala 150	
tac acc cgc tac atg ggg gat ctc tcg ggg ggc cag gtg ctg aag aag	591
Tyr Thr Arg Tyr Met 155 Gly Asp Leu Ser Gly Gly Gln Val Leu Lys Lys 160 165	
gtg gcc cag cga gca ctg aaa ctc ccc agc aca ggg gaa ggg acc cag	639
Val Ala Gln Arg Ala Leu Lys Leu Pro Ser Thr Gly Glu Gly Thr Gln 170 175 180 185	
ttc tac ctg ttt gag aat gtg gac aat gcc cag cag ttc aag cag ctc	687
Phe Tyr Leu Phe 190 Glu Asn Val Asp Asn Ala Gln Gln Phe Lys Gln Leu 195 200	
tac cgg gcc agg atg aac gcc ctg gac ctg aac atg aag acc aaa gag	735
Tyr Arg Ala Arg Met 205 Asn Ala Leu Asp Leu Asn Met Lys Thr Lys Glu 210 215	
agg atc gtg gag gcc aac aag gct ttt gag tat aac atg cag ata ttc	783
Arg Ile Val Glu Ala Asn Lys Ala Phe Glu Tyr Asn Met Gln Ile Phe 220 225 230	
aat gaa ctg gac cag gcc ggc tcc aca ctg gcc aga gag acc ttg gag	831
Asn Glu Leu Asp Gln Ala Gly Ser Thr Leu Ala Arg Glu Thr Leu Glu 235 240 245	
gat ggg ttc cct gta cac gat ggg aaa gga gac atg cgt aaa tgc cct	879
Asp Gly Phe Pro Val His Asp Gly Lys Gly Asp Met Arg Lys Cys Pro 250 255 260 265	
ttc tac gct gct gaa caa gac aaa ggg ctg gag ggc agc ctg tcc ctt	927
Phe Tyr Ala Ala Glu Gln Asp Lys Gly Leu Glu Gly Ser Leu Ser Leu 270 275 280	
ccg aca agc tat gct gtg ctg agg aag ccc agc ctc cag ttc atc ctg	975
Pro Thr Ser Tyr Ala Val Leu Arg Lys Pro Ser Leu Gln Phe Ile Leu 285 290 295	
gcc gct ggt gtg gcc cta gct gct gga ctc ttg gcc tgg tac tac atg	1023
Ala Ala Gly Val Ala Leu Ala Ala Gly Leu Leu Ala Trp Tyr Tyr Met 300 305 310	
tgaagcacc atcatgccac accggtaccc tccctcccgac tgaccactgg cctaccctt	1083
tctccagccc tgactaaact accacctcag gtgacttttt aaaaaatgct gggtttaaga	1143
aaggcaacca ataaaagaga tgctagagcc tcgtctgaca gcacacctc tatgggcat	1203
attccgcact gggcacaggc cgtcaccctg ggagcagtcg gcacagtgca gcaagcctgg	1263
cccccgaccc agctctactc caggcttcca cacttctggg ccctaggctg cttccggtag	1323
tcctgtttt tgcagtacat ggggtgactat ctccctgtt ggaggtgagt ggctgttaag	1383
tccaagctgt gcgagggggc cttgctggat gctgctgtac aacttctggg cctctcttgg	1443
accctgggag tgaggggtggg tgtgggtgga agcctcagag gccttgggag ctcacccctc	1503
tcaccagaa tccctctaac ccttgggtgc ggtttgctca gcccagctt atctcctcct	1563
ccgctgtgt aaatgctcca gcactcaata aagtgggctt tgcaagctaa aaaaaaaaaa	1623
aaaa	1627

<210> 4
 <211> 313
 <212> PRT

<213> Homo sapiens

<400> 4

Met Ser Ala Glu Val Glu Thr Ser Glu Gly Val Asp Glu Ser Glu Lys
 1 5 10 15
 Lys Asn Ser Gly Ala Leu Glu Lys Glu Asn Gln Met Arg Met Ala Asp
 20 25 30
 Leu Ser Glu Leu Leu Lys Glu Gly Thr Lys Glu Ala His Asp Arg Ala
 35 40 45
 Glu Asn Thr Gln Phe Val Lys Asp Phe Leu Lys Gly Asn Ile Lys Lys
 50 55 60
 Glu Leu Phe Lys Leu Ala Thr Thr Ala Leu Tyr Phe Thr Tyr Ser Ala
 65 70 75 80
 Leu Glu Glu Glu Met Glu Arg Asn Lys Asp His Pro Ala Phe Ala Pro
 85 90 95
 Leu Tyr Phe Pro Met Glu Leu His Arg Lys Glu Ala Leu Thr Lys Asp
 100 105 110
 Met Glu Tyr Phe Phe Gly Glu Asn Trp Glu Glu Gln Val Gln Cys Pro
 115 120 125
 Lys Ala Ala Gln Lys Tyr Val Glu Arg Ile His Tyr Ile Gly Gln Asn
 130 135 140
 Glu Pro Glu Leu Leu Val Ala His Ala Tyr Thr Arg Tyr Met Gly Asp
 145 150 155 160
 Leu Ser Gly Gly Gln Val Leu Lys Lys Val Ala Gln Arg Ala Leu Lys
 165 170 175
 Leu Pro Ser Thr Gly Glu Gly Thr Gln Phe Tyr Leu Phe Glu Asn Val
 180 185 190
 Asp Asn Ala Gln Gln Phe Lys Gln Leu Tyr Arg Ala Arg Met Asn Ala
 195 200 205
 Leu Asp Leu Asn Met Lys Thr Lys Glu Arg Ile Val Glu Ala Asn Lys
 210 215 220
 Ala Phe Glu Tyr Asn Met Gln Ile Phe Asn Glu Leu Asp Gln Ala Gly
 225 230 235 240
 Ser Thr Leu Ala Arg Glu Thr Leu Glu Asp Gly Phe Pro Val His Asp
 245 250 255
 Gly Lys Gly Asp Met Arg Lys Cys Pro Phe Tyr Ala Ala Glu Gln Asp
 260 265 270
 Lys Gly Leu Glu Gly Ser Leu Ser Leu Pro Thr Ser Tyr Ala Val Leu
 275 280 285
 Arg Lys Pro Ser Leu Gln Phe Ile Leu Ala Ala Gly Val Ala Leu Ala
 290 295 300
 Ala Gly Leu Leu Ala Trp Tyr Tyr Met
 305 310

<210> 5

<211> 2225

<212> DNA

<213> Rattus rattus

<220>

<221> CDS

<222> (1062)...(1931)

<400> 5

tttcaggggat	ttttgcgatt	cctctctgtgta	gacttctact	tggtctctctaa	gggagttctt	60
catgtcttttc	ttgaagtcac	ccagcatcat	gatcaaatat	gattttgaaa	ctagatcttg	120
cttttctgggt	gtgtttggat	attccatggt	tggtttgggtg	ggagaattgg	gctccgatga	180
tggtcatgtag	tcttggtttc	tggtgcttggtg	tttcctgcgc	ttgcctctcg	ccatcagatt	240
atctctagtgt	ttactttgtt	ctgctatttc	tgacagtggc	tagactgtcc	tataagcctg	300
tgtgtcagga	gtgctgtaga	cctttttttcc	tctcttttcag	tcagttatgg	gacagagtgt	360
tctgctttttg	ggcgtgtagt	ttttcctctc	tacaggtctt	cagctgttcc	tgtgggcctg	420
tgtcttgagt	tcaccaggca	gctttcttgc	agcagaaaaa	ttggtcatac	ctgtgatcct	480
gaggctcaag	ttegtctgtg	gggtgctgtc	caggggctct	ctgcagcggg	cacaaccagg	540
aagacctgtg	cggcccttc	cggagcttca	gtgcaccagg	gttccagatg	gcctttggcg	600
ttttcctctg	cggctccgaga	tgtatgtaca	gagagcagtc	tcttctgggt	tcccaggcct	660
gtctgcctct	ctgaagggtc	agctctccct	cccacgggat	ttgggtgcag	agaactgttt	720
atccgggtctg	tttctttcag	gttccgggtg	tgtctcaggc	aggtgtcgtt	cctgcgcctt	780

ccccatggg accagaggcc ttatacagtt tcctcttggg ccagggatgt gggcaggggt	840
gagcagtgtt ggtggtctct tccgtctgca gcctcaggag tgccacctga ccaggcgggt	900
gggtctctct ctgagaattt cattttttaa tcattcatta aaatgtcatg acttgatgtc	960
ctgctgtccg tctcacgccc tcagctgtaa cagtgccgag ggagtcactg aagaagagac	1020
tgaatgacca gagtatgggc agcacagaca actcaacaaa a atg tct tca gag gtg	1076
Met Ser Ser Glu Val	
1 5	
gag act gcg gag gcc gta gat gag tca gag aag aac tct atg gca tca	1124
Glu Thr Ala Glu Ala Val Asp Glu Ser Glu Lys Asn Ser Met Ala Ser	
10 15 20	
gag aag gaa aac cat tcc aaa ata gca gac ttt tct gat ctt ctg aag	1172
Glu Lys Glu Asn His Ser Lys Ile Ala Asp Phe Ser Asp Leu Leu Lys	
25 30 35	
gaa ggg aca aag gaa gca gat gac cgg gca gaa aat acc cag ttt gtc	1220
Glu Gly Thr Lys Glu Ala Asp Asp Arg Ala Glu Asn Thr Gln Phe Val	
40 45 50	
aaa gac ttc ttg aaa gga aac att aag aag gag cta ttt aag ctg gcc	1268
Lys Asp Phe Leu Lys Gly Asn Ile Lys Lys Glu Leu Phe Lys Leu Ala	
55 60 65	
acc act gca ctt tca tac tca gcc cct gag gag gaa atg gat tca ctg	1316
Thr Thr Ala Leu Ser Tyr Ser Ala Pro Glu Glu Met Asp Ser Leu	
70 75 80 85	
acc aag gac atg gag tac ttc ttt ggt gaa aac tgg gag gaa aaa gtg	1364
Thr Lys Asp Met Glu Tyr Phe Phe Gly Glu Asn Trp Glu Glu Lys Val	
90 95 100	
aag tgc tct gaa gct gcc cag acg tat gtg gat cag att cac tat gta	1412
Lys Cys Ser Glu Ala Ala Gln Thr Tyr Val Asp Gln Ile His Tyr Val	
105 110 115	
ggg caa aat gag cca gag cat ctg gtg gcc cat act tac tct act tac	1460
Gly Gln Asn Glu Pro Glu His Leu Val Ala His Thr Tyr Ser Thr Tyr	
120 125 130	
atg ggg gga aac ctt tca ggg gac cag gta ctg aag aag gag acc cag	1508
Met Gly Gly Asn Leu Ser Gly Asp Gln Val Leu Lys Lys Glu Thr Gln	
135 140 145	
ccg gtc ccc ttc act agg gaa ggg act cag ttc tac ctg ttt gag cat	1556
Pro Val Pro Phe Thr Arg Glu Gly Thr Gln Phe Tyr Leu Phe Glu His	
150 155 160 165	
gta gac aat gct aag caa ttc aag cta ttc tac tgc gct aga ttg aat	1604
Val Asp Asn Ala Lys Gln Phe Lys Leu Phe Tyr Cys Ala Arg Leu Asn	
170 175 180	
gcc ttg gac ctg aat ttg aag acc aaa gag agg att gtg gag gaa gcc	1652
Ala Leu Asp Leu Asn Leu Lys Thr Lys Glu Arg Ile Val Glu Glu Ala	
185 190 195	
acc aaa gcc ttt gaa tat aat atg cag ata ttc agt gaa ctg gac cag	1700
Thr Lys Ala Phe Glu Tyr Asn Met Gln Ile Phe Ser Glu Leu Asp Gln	
200 205 210	
gca ggc tcc ata cca gta aga gaa acc cta aag aat ggg ctc tca ata	1748
Ala Gly Ser Ile Pro Val Arg Glu Thr Leu Lys Asn Gly Leu Ser Ile	
215 220 225	
ctt gat ggg aag gga ggt gta tgc aaa tgt ccc ttt aat gct gct cag	1796
Leu Asp Gly Lys Gly Gly Val Cys Lys Cys Pro Phe Asn Ala Ala Gln	
230 235 240 245	

cca gac aaa ggt acc ctg gga ggc agc aac tgc cct ttc cag atg tcc 1844
Pro Asp Lys Gly Thr Leu Gly Gly Ser Asn Cys Pro Phe Gln Met Ser 260
250 255

atg gcc ttg ctg agg aag cct aac ttg cag ctc att cta gtt gcc agt 1892
Met Ala Leu Leu Arg Lys Pro Asn Leu Gln Leu Ile Leu Val Ala Ser 275
265 270

atg gcc ttg gta gct gga ctt tta gcc tgg tac tac atg tgaagggcct 1941
Met Ala Leu Val Ala Gly Leu Leu Ala Trp Tyr Tyr Met 290
280 285

gtcaagttgt ttgcataccta tctcaacatc ctaccacttg ttccttcccc acctccacct 2001
ctgcctagaa ctaccacctc aggtgacatt tttaatgttg ggtttgagaa aatgagcaac 2061
caataaaaaga cagaccctag aaaaaagtca tgacttaagt ggcacgggga cacctaaagt 2121
cacactttgt gcttcagaca tactttcttt ctctatttca acactgaatt cggaagtaa 2181
cctactacta ttaataataa atgctacaca atgcataata aaaa 2225

<210> 6
<211> 290
<212> PRT
<213> Rattus rattus

<400> 6
Met Ser Ser Glu Val Glu Thr Ala Glu Ala Val Asp Glu Ser Glu Lys 15
1 5 10
Asn Ser Met Ala Ser Glu Lys Glu Asn His Ser Lys Ile Ala Asp Phe 30
20 25
Ser Asp Leu Lys Glu Gly Thr Lys Glu Ala Asp Asp Arg Ala Glu 45
35 40
Asn Thr Gln Phe Val Lys Asp Phe Leu Lys Gly Asn Ile Lys Lys Glu 60
50 55
Leu Phe Lys Leu Ala Thr Thr Ala Leu Ser Tyr Ser Ala Pro Glu Glu 80
65 70
Glu Met Asp Ser Leu Thr Lys Asp Met Glu Tyr Phe Phe Gly Glu Asn 95
85 90
Trp Glu Glu Lys Val Lys Cys Ser Glu Ala Ala Gln Thr Tyr Val Asp 110
100 105
Gln Ile His Tyr Val Gly Gln Asn Glu Pro Glu His Leu Val Ala His 125
115 120
Thr Tyr Ser Thr Tyr Met Gly Gly Asn Leu Ser Gly Asp Gln Val Leu 140
130 135
Lys Lys Glu Thr Gln Pro Val Pro Phe Thr Arg Glu Gly Thr Gln Phe 160
145 150
Tyr Leu Phe Glu His Val Asp Asn Ala Lys Gln Phe Lys Leu Phe Tyr 175
165 170
Cys Ala Arg Leu Asn Ala Leu Asp Leu Asn Leu Lys Thr Lys Glu Arg 190
180 185
Ile Val Glu Glu Ala Thr Lys Ala Phe Glu Tyr Asn Met Gln Ile Phe 205
195 200
Ser Glu Leu Asp Gln Ala Gly Ser Ile Pro Val Arg Glu Thr Leu Lys 220
210 215
Asn Gly Leu Ser Ile Leu Asp Gly Lys Gly Gly Val Cys Lys Cys Pro 240
225 230 235
Phe Asn Ala Ala Gln Pro Asp Lys Gly Thr Leu Gly Gly Ser Asn Cys 255
245 250
Pro Phe Gln Met Ser Met Ala Leu Leu Arg Lys Pro Asn Leu Gln Leu 270
260 265
Ile Leu Val Ala Ser Met Ala Leu Val Ala Gly Leu Leu Ala Trp Tyr 285
275 280
Tyr Met 290

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 99/19823

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K38/44 A61K48/00 A61P9/10 //A61K38/18,A61K31/555

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ABRAHAM, N. G. (1) ET AL: "Transfection of the human heme oxygenase gene into rabbit coronary microvessel endothelial cells: Protective effect against heme and hemoglobin toxicity." PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, (1995) VOL. 92, NO. 15, PP. 6798-6802. , XP002100374 the whole document	1-12
X	WO 98 08566 A (WISCONSIN MED COLLEGE INC ;UNIV DUKE (US)) 5 March 1998 (1998-03-05) page 4, line 23 -page 6, line 12 page 7, line 24 -page 8, line 2 page 9, line 19 -page 10, line 18 page 55, line 2 - line 30 page 57, line 22 - line 28 -/-	14-25

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

23 February 2000

Date of mailing of the international search report

09/03/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Stein, A

INTERNATIONAL SEARCH REPORT

Application No

PCT/US 99/19823

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 97 36615 A (HARVARD COLLEGE) 9 October 1997 (1997-10-09) page 2, line 11 - line 36 page 7, line 16 -page 8, line 13 page 16, line 18 -page 17, line 36 page 21, line 14 -page 22, line 8 claims 1-6 ---	14-23
A	MAULIK N ET AL: "Nitric oxide/carbon monoxide. A molecular switch for myocardial preservation during ischemia." CIRCULATION, (1996 NOV 1) 94 (9 SUPPL) II398-406. , XP000876907 the whole document ---	1-13
A	ABRAHAM, NADER G. (1): "Manipulation of heme oxygenase expression by gene transfer and metals: Implications in cell injury and repair." JOURNAL OF NEUROCHEMISTRY, (1998) VOL. 70, NO. SUPPL. 1, PP. S45. MEETING INFO.: 29TH ANNUAL MEETING OF THE AMERICAN SOCIETY FOR NEUROCHEMISTRY DENVER, COLORADO, USA MARCH 7-11, 1998 AMERICAN SOCIETY FOR NEUROCHEMISTRY. , XP000876935 the whole document ---	1-13
A	LONG, XILIN ET AL: "Hypoxia-induced expression of heme oxygenase gene expression in cultured neonatal rat cardiac myocytes." CIRCULATION, (1995) VOL. 92, NO. 8 SUPPL., PP. I653-I654, XP000876926 the whole document ---	1-13
A	HOSHIDA, SHIRO ET AL: "Heme oxygenase -1 as a culture shock protein in rat neonatal cardiomyocytes." JOURNAL OF MOLECULAR AND CELLULAR CARDIOLOGY, (1994) VOL. 26, NO. 11, PP. CCXII, XP000876927 the whole document ---	1-13
A	BORGER DR: "Induction of HSP 32 gene in hypoxic cardiomyocytes is attenuated by treatment with N-acetyl-L-cysteine" AMERICAN JOURNAL OF PHYSIOLOGY, vol. 274, no. 3 Pt 2, March 1998 (1998-03), pages H965-73, XP002131421 the whole document ---	1-14
	---	1-14

-/--

INTERNATIONAL SEARCH REPORT

Application No
PCT/US 99/19823

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	MORITA T ET AL: "Carbon monoxide controls the proliferation of hypoxic vascular smooth muscle cells" JOURNAL OF BIOLOGICAL CHEMISTRY, (26 DEC 1997) VOL. 272, NO. 52, PP. 32804-32809, XP002131422 the whole document	24,25
P,X	SOARES M P ET AL: "Expression of heme oxygenase -1 can determine cardiac xenograft survival." NATURE MEDICINE, (1998 SEP) 4 (9) 1073-7., XP002131423 the whole document	13

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/ 19823

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 1-8, 14-25 are directed to a method of treatment of the human/animal the search has been carried out and based on the alleged effects of the compound /composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/19823

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
---	---------------------	----------------------------	---------------------

WO 9808566 A	05-03-1998	AU 4054297 A EP 0963219 A	19-03-1998 15-12-1999
--------------	------------	------------------------------	--------------------------

WO 9736615 A	09-10-1997	US 5888982 A	30-03-1999
--------------	------------	--------------	------------